

Abstract

We suggest a so-called Dirac type tensor equation with nonabelian gauge symmetry on pseudo-Riemannian space. This equation reproduce some of the properties of spinor Dirac equation. A geometrical interpretation of results in terms of Riemannian geometry is given.

Dirac-type tensor equations with non-Abelian gauge symmetries on pseudo-Riemannian space

N.G.Marchuk *

February 7, 2008

PACS: 04.20Cv, 04.62, 11.15, 12.10

Steklov Mathematical Institute, Gubkina st.8, Moscow 119991, Russia
nmarchuk@mi.ras.ru, www.orc.ru/~nmarchuk

Contents

1	Basic definitions.	3
2	Lie algebras of anti-Hermitian differential forms.	13
3	Main equations.	16
4	Nonabelian charge conservation laws.	18
5	Unitary and Spin gauge symmetries.	20
6	Lagrangians.	21
7	A geometrical interpretation of results.	23

*Research supported by the Russian Foundation for Basic Research grants 00-01-00224, 00-15-96073.

8	Space dimensions $n = 2, 3$.	25
9	Addendum.	26

In the previous paper [3], developing results of P. Dirac [4], D. D. Ivanenko, L. D. Landau [5], E. Kähler [6], F. Gürsey [7], D. Hestenes [8], [9], we consider the so-called Dirac type tensor equation which reproduce some of the properties of the spinor Dirac equation for an electron. Now we generalize the Dirac type tensor equation in two ways.

Firstly we present the Dirac type tensor equation with nonabelian gauge symmetry. We use unitary gauge Lie groups that are subgroups of the group $U(4)$, in particular $U(1)$, $U(1) \times SU(2)$, $SU(3)$ – main groups of the Standard Model. Note that in case of the gauge group $SU(3)$ the Dirac type tensor equation has 16 complex valued components of wave function. But the Dirac equation for chromospinor has only 12 complex valued components. Evidently these two systems of equations are not coincide.

Secondly we generalize all results on pseudo-Riemannian space (in [3] equations are considered in Minkowski space). A method of generalization was suggested in [2]. A key role in this method play the tensor B_μ . Now we have found explicit formulas for B_μ via the components of metric tensor $g_{\mu\nu}$ and its first derivatives. In the section 7 we give an interpretation of our results in terms of Riemannian geometry.

1 Basic definitions.

More details about the following definitions can be found in [2].

1. Let $\mathcal{M}$ be a four dimensional differentiable manifolds with a local system of coordinates x^μ . Greek indices run over (1,2,3,4). Summation convention over repeating indices is assumed. Suppose that there is a smooth twice covariant tensor field (metric tensor) with components $g_{\mu\nu} = g_{\mu\nu}(x)$, $x \in \mathcal{M}$ such that

- $g_{\mu\nu} = g_{\nu\mu}$;
- $g = \det \|g_{\mu\nu}\| < 0$;
- The signature of the matrix $\|g_{\mu\nu}\|$ is equal to -2 .

The matrix $\|g^{\mu\nu}\|$ composed from contravariant components of the metric tensor is the inverse matrix to $\|g_{\mu\nu}\|$. The full set of $\{\mathcal{M}, g_{\mu\nu}\}$ is called a *pseudo-Riemannian space* and denoted by $\mathcal{V}$.

2. Let Λ_k be the sets of exterior differential forms of rank $k = 0, 1, 2, 3, 4$ on $\mathcal{V}$ (covariant antisymmetric tensor fields) and

$$\Lambda = \Lambda_0 \oplus \dots \oplus \Lambda_4 = \Lambda_{\text{ev}} \oplus \Lambda_{\text{od}}, \quad \Lambda_{\text{ev}} = \Lambda_0 \oplus \Lambda_2 \oplus \Lambda_4, \quad \Lambda_{\text{od}} = \Lambda_1 \oplus \Lambda_3.$$

Elements of Λ are called (nonhomogeneous) *differential forms* and elements of Λ_k are called *k-forms* or differential forms of rank k . Elements of Λ_{ev} and Λ_{od} are called *even* and *odd* forms respectively. The set of smooth scalar functions on $\mathcal{V}$ (invariants) is identified with the set of 0-forms Λ_0 . A k -form $U \in \Lambda_k$ can be written as

$$U = \frac{1}{k!} u_{\nu_1 \dots \nu_k} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_k} = \sum_{\mu_1 < \dots < \mu_k} u_{\mu_1 \dots \mu_k} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_k}, \quad (1)$$

where the smooth functions $u_{\nu_1 \dots \nu_k} = u_{\nu_1 \dots \nu_k}(x)$ are real valued components of a covariant antisymmetric ($u_{\nu_1 \dots \nu_k} = u_{[\nu_1 \dots \nu_k]}$) tensor field.

Let $\Lambda_k^{\mathcal{C}}, \Lambda^{\mathcal{C}}, \Lambda_{\text{ev}}^{\mathcal{C}}, \Lambda_{\text{od}}^{\mathcal{C}}$ be corresponding sets of complex valued differential forms and $\mathcal{K}$ be a field of complex numbers $\mathcal{C}$ or real numbers $\mathcal{R}$ ($\Lambda^{\mathcal{R}} = \Lambda$). Differential forms from $\Lambda^{\mathcal{K}}$ can be written as linear combinations of the 16 basis differential forms

$$1, dx^{\mu}, dx^{\mu_1} \wedge dx^{\mu_2}, \dots, dx^0 \wedge \dots \wedge dx^3, \quad \mu_1 < \mu_2 < \dots \quad (2)$$

with coefficients from $C^\infty(\mathcal{V}, \mathcal{K})$ (smooth functions that map $\mathcal{V}$ into $\mathcal{K}$). The exterior product of differential forms is defined in the usual way. If $U \in \Lambda_r^{\mathcal{K}}, V \in \Lambda_s^{\mathcal{K}}$, then

$$U \wedge V = (-1)^{rs} V \wedge U \in \Lambda_{r+s}^{\mathcal{K}}.$$

3. Consider the Hodge star operator $\star : \Lambda_k^{\mathcal{K}} \rightarrow \Lambda_{4-k}^{\mathcal{K}}$. If $U \in \Lambda_k$ has the form (1), then

$$\star U = \frac{1}{k!(4-k)!} \sqrt{-g} \varepsilon_{\mu_1 \dots \mu_4} u^{\mu_1 \dots \mu_k} dx^{\mu_{k+1}} \wedge \dots \wedge dx^{\mu_4},$$

where $u^{\mu_1 \dots \mu_k} = g^{\mu_1 \nu_1} \dots g^{\mu_k \nu_k} u_{\nu_1 \dots \nu_k}$, $\varepsilon_{\mu_1 \dots \mu_4}$ is the sign of the permutation $(\mu_1 \dots \mu_4)$, and $\varepsilon_{0123} = 1$. It is easy to prove that for $U \in \Lambda_k^{\mathcal{K}}$

$$\star(\star U) = (-1)^{k+1} U.$$

The form $\star U$ is a covariant antisymmetric tensor with respect to changes of coordinates with positive Jacobian.

4. Let us define the *volume form*

$$\ell = \sqrt{-g} dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3 = \frac{\sqrt{-g}}{4!} \epsilon_{\mu_1 \dots \mu_4} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_4}.$$

We have

$$(\ell)^2 = -1, \quad \overset{k}{U} \ell = (-1)^k \ell \overset{k}{U}, \quad \text{for} \quad \overset{k}{U} \in \Lambda_k$$

that means ℓ commutes with all even forms and anticommutes with all odd forms with respect to Clifford product (see below).

5. Further on we consider the bilinear operator $\text{Com} : \Lambda_2 \times \Lambda_2 \rightarrow \Lambda_2$ such that

$$\begin{aligned} \text{Com}\left(\frac{1}{2}a_{\mu_1\mu_2}dx^{\mu_1} \wedge dx^{\mu_2}, \frac{1}{2}b_{\nu_1\nu_2}dx^{\nu_1} \wedge dx^{\nu_2}\right) &= \frac{1}{2}a_{\mu_1\mu_2}b_{\nu_1\nu_2}(-g^{\mu_1\nu_1}dx^{\mu_2} \wedge dx^{\nu_2} \\ &- g^{\mu_2\nu_2}dx^{\mu_1} \wedge dx^{\nu_1} + g^{\mu_1\nu_2}dx^{\mu_2} \wedge dx^{\nu_1} + g^{\mu_2\nu_1}dx^{\mu_1} \wedge dx^{\nu_2}), \end{aligned}$$

where $a_{\mu_1\mu_2} = a_{[\mu_1\mu_2]}$, $b_{\nu_1\nu_2} = b_{[\nu_1\nu_2]}$. Evidently, $\text{Com}(U, V) = -\text{Com}(V, U)$.

6. Now we define the Clifford product of differential forms with the aid of the following formulas:

$$\begin{aligned} \overset{0}{U} \overset{k}{V} &= \overset{k}{V} \overset{0}{U} = \overset{0}{U} \wedge \overset{k}{V} = \overset{k}{V} \wedge \overset{0}{U}, \\ \overset{1}{U} \overset{k}{V} &= \overset{1}{U} \wedge \overset{k}{V} - \star(\overset{1}{U} \wedge \star \overset{k}{V}), \\ \overset{k}{U} \overset{1}{V} &= \overset{k}{U} \wedge \overset{1}{V} + \star(\star \overset{k}{U} \wedge \overset{1}{V}), \\ \overset{2}{U} \overset{2}{V} &= \overset{2}{U} \wedge \overset{2}{V} + \star(\overset{2}{U} \wedge \star \overset{2}{V}) + \frac{1}{2}\text{Com}(\overset{2}{U}, \overset{2}{V}), \\ \overset{2}{U} \overset{3}{V} &= \star \overset{2}{U} \wedge \star \overset{3}{V} - \star(\overset{2}{U} \wedge \star \overset{3}{V}), \\ \overset{2}{U} \overset{4}{V} &= \star \overset{2}{U} \wedge \star \overset{4}{V}, \\ \overset{3}{U} \overset{2}{V} &= -\star \overset{3}{U} \wedge \star \overset{2}{V} - \star(\star \overset{3}{U} \wedge \overset{2}{V}), \\ \overset{3}{U} \overset{3}{V} &= \star \overset{3}{U} \wedge \star \overset{3}{V} + \star(\overset{3}{U} \wedge \star \overset{3}{V}), \\ \overset{3}{U} \overset{4}{V} &= \star \overset{3}{U} \wedge \star \overset{4}{V}, \\ \overset{4}{U} \overset{2}{V} &= \star \overset{4}{U} \wedge \star \overset{2}{V}, \end{aligned}$$

$$\begin{aligned}\overset{4}{U}\overset{3}{V} &= -\star \overset{4}{U} \wedge \star \overset{3}{V}, \\ \overset{4}{U}\overset{4}{V} &= -\star \overset{4}{U} \wedge \star \overset{4}{V},\end{aligned}$$

where ranks of differential forms are denoted as $\overset{k}{U} \in \Lambda_k^\mathcal{K}$ and $k = 0, 1, 2, 3, 4$. From this definition we may obtain properties of the Clifford product of differential forms.

- If $U, V \in \Lambda^\mathcal{K}$, then $UV \in \Lambda^\mathcal{K}$.
- The axioms of associativity and distributivity are satisfied for the Clifford product.
- $dx^\mu dx^\nu = dx^\mu \wedge dx^\nu + g^{\mu\nu}$, $dx^\mu dx^\nu + dx^\nu dx^\mu = 2g^{\mu\nu}$.
- $\text{Com}(U, V) = UV - VU$ for $U, V \in \Lambda_2^\mathcal{K}$.

7. Let us define the trace of differential forms as the linear operation $\text{Tr} : \Lambda^\mathcal{K} \rightarrow \Lambda_0^\mathcal{K}$ such that

$$\text{Tr}(1) = 1, \quad \text{Tr}(dx^{\mu_1} \wedge \dots \wedge dx^{\mu_k}) = 0 \quad \text{for } k = 1, 2, 3, 4.$$

The reader can easily prove that

$$\text{Tr}(UV - VU) = 0, \quad \text{Tr}(V^{-1}UV) = \text{Tr } U, \quad U, V \in \Lambda^\mathcal{K}.$$

In the second relation V is an invertible exterior form with respect to (w.r.t.) Clifford product.

8. Let us define the involution $*$: $\Lambda_k^\mathcal{K} \rightarrow \Lambda_k^\mathcal{K}$. By definition, put

$$U^* = (-1)^{\frac{k(k-1)}{2}} \check{U}, \quad U \in \Lambda_k^\mathcal{K},$$

where $\check{U}$ is the differential form with complex conjugated components (if $\mathcal{K} = \mathcal{R}$, then $\check{U} = U$). It is readily seen that

$$U^{**} = U, \quad (UV)^* = V^* U^*, \quad U, V \in \Lambda^\mathcal{K}.$$

9. Now we can define the spinor group

$$\text{Spin}(\mathcal{V}) = \{S \in \Lambda_{\text{ev}} : S^* S = 1\}.$$

Note that this definition is valid only for space dimensions $n < 6$ (see [2]).

10. Let

$$u_{\mu_1 \dots \mu_k \nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r}(x) = u_{[\mu_1 \dots \mu_k] \nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r}(x), \quad x \in \mathcal{V}$$

be components of a tensor field of rank $(r, s+k)$ antisymmetric with respect to the first k covariant indices. One may consider the following objects:

$$U_{\nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r} = \frac{1}{k!} u_{\mu_1 \dots \mu_k \nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_k} \quad (3)$$

which are formally written as k -forms. Under a change of coordinates $(x) \rightarrow (\tilde{x})$ the values (3) transform as components of tensor field of rank (r, s)

$$\tilde{U}_{\beta_1 \dots \beta_s}^{\alpha_1 \dots \alpha_r} = q_{\beta_1}^{\nu_1} \dots q_{\beta_s}^{\nu_s} p_{\lambda_1}^{\alpha_1} \dots p_{\lambda_r}^{\alpha_r} U_{\nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r}, \quad q_{\beta}^{\nu} = \frac{\partial x^{\nu}}{\partial \tilde{x}^{\beta}}, \quad p_{\lambda}^{\alpha} = \frac{\partial \tilde{x}^{\alpha}}{\partial x^{\lambda}}. \quad (4)$$

The objects (3) are called tensors of rank (r, s) with values in $\Lambda_k^{\mathcal{K}}$. We write this as

$$U_{\nu_1 \dots \nu_s}^{\lambda_1 \dots \lambda_r} \in \Lambda_k^{\mathcal{K}} \top_s^r.$$

Elements of $\Lambda_0^{\mathcal{K}} \top_s^r$ are ordinary tensor fields of rank (r, s) on $\mathcal{V}$. Let us define the Clifford product of the elements $U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} \in \Lambda_s^{\mathcal{K}} \top_s^r$ and $V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p} \in \Lambda_q^{\mathcal{K}} \top_q^p$ as the tensor field from $\Lambda_{s+q}^{\mathcal{K}} \top_{s+q}^{r+p}$ of the form

$$W_{\nu_1 \dots \nu_s \beta_1 \dots \beta_q}^{\mu_1 \dots \mu_r \alpha_1 \dots \alpha_p} = U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p},$$

where at right hand side there is the Clifford product of differential forms (the indices $\mu_1, \dots, \mu_r, \alpha_1, \dots, \alpha_p, \nu_1, \dots, \nu_s, \beta_1, \dots, \beta_q$ are fixed). In particular, it follows from this definition that if $U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} \in \Lambda_0^{\mathcal{K}} \top_s^r$ and $V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p} \in \Lambda_0^{\mathcal{K}} \top_q^p$, then the Clifford product of these elements is identified with the tensor product.

Note that $dx^{\mu} = \delta_{\nu}^{\mu} dx^{\nu}$, where δ_{ν}^{μ} is the Kronecker tensor ($\delta_{\nu}^{\mu} = 0$ for $\mu \neq \nu$ and $\delta_{\mu}^{\mu} = 1$). Hence, $dx^{\mu} \in \Lambda_1^{\mathcal{K}} \top_1^1$.

11. Let us define the *Upsilon derivatives* Υ_{μ} which act on tensors from $\Lambda_s^{\mathcal{K}} \top_s^r$ by the following rules:

a) If $t_{\nu_1 \dots \nu_s}^{\epsilon_1 \dots \epsilon_r} \in \Lambda_0^{\mathcal{K}} \top_s^r$, then

$$\Upsilon_{\mu} t_{\nu_1 \dots \nu_s}^{\epsilon_1 \dots \epsilon_r} = \partial_{\mu} t_{\nu_1 \dots \nu_s}^{\epsilon_1 \dots \epsilon_r}.$$

b) $\Upsilon_{\mu} dx^{\nu} = -\Gamma^{\nu}_{\mu\lambda} dx^{\lambda}$, where $\Gamma^{\mu}_{\mu\lambda} = \Gamma^{\mu}_{\lambda\mu}$ are Christoffel symbols (Levi-Civita connectedness components).

c) If $U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} \in \Lambda^{\mathcal{K}} \mathbb{T}_s^r$, $V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p} \in \Lambda^{\mathcal{K}} \mathbb{T}_q^p$, then

$$\Upsilon_\lambda(U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p}) = (\Upsilon_\lambda U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r}) V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p} + U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} (\Upsilon_\lambda V_{\beta_1 \dots \beta_q}^{\alpha_1 \dots \alpha_p}).$$

d) If $U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r}, V_{\beta_1 \dots \beta_s}^{\alpha_1 \dots \alpha_r} \in \Lambda^{\mathcal{K}} \mathbb{T}_s^r$, then

$$\Upsilon_\lambda(U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} + V_{\beta_1 \dots \beta_s}^{\alpha_1 \dots \alpha_r}) = \Upsilon_\lambda U_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_r} + \Upsilon_\lambda V_{\beta_1 \dots \beta_s}^{\alpha_1 \dots \alpha_r}.$$

With the aid of these rules it is easy to calculate how operators Υ_μ act on arbitrary tensor from $\Lambda^{\mathcal{K}} \mathbb{T}_s^r$.

From the formula $\Upsilon_\mu dx^\lambda = -\Gamma^\lambda_{\mu\nu} dx^\nu$ we get

$$(\Upsilon_\mu \Upsilon_\nu - \Upsilon_\nu \Upsilon_\mu) dx^\lambda = -R^\lambda_{\rho\mu\nu} dx^\rho, \quad (5)$$

where

$$R^\kappa_{\lambda\mu\nu} = \partial_\mu \Gamma^\kappa_{\nu\lambda} - \partial_\nu \Gamma^\kappa_{\mu\lambda} + \Gamma^\kappa_{\mu\eta} \Gamma^\eta_{\nu\lambda} - \Gamma^\kappa_{\nu\eta} \Gamma^\eta_{\mu\lambda} \quad (6)$$

is rank (1,3) tensor, known as the curvature tensor (or Riemannian tensor).

12. Consider the antisymmetric tensor from $\Lambda_2 \mathbb{T}_2$

$$C_{\mu\nu} = \frac{1}{2} R_{\alpha\beta\mu\nu} dx^\alpha \wedge dx^\beta,$$

where $R_{\alpha\beta\mu\nu} = g_{\alpha\lambda} R^\lambda_{\beta\mu\nu}$. Let $B_\mu \in \Lambda_2 \mathbb{T}_1$ be such that

$$\Upsilon_\mu B_\nu - \Upsilon_\nu B_\mu - [B_\mu, B_\nu] = \frac{1}{2} C_{\mu\nu}. \quad (7)$$

Existence of B_μ is considered in the Theorem 2.

13. Let us define the linear differential operators

$$\mathcal{D}_\mu = \Upsilon_\mu - [B_\mu, \cdot].$$

Theorem 1. *The operators $\mathcal{D}_\mu$ satisfy Leibniz rule*

$$\mathcal{D}_\mu(UV) = (\mathcal{D}_\mu U)V + U\mathcal{D}_\mu V \quad \text{for } U \in \Lambda^{\mathcal{K}} \mathbb{T}_s^r, \quad V \in \Lambda^{\mathcal{K}} \mathbb{T}_q^p$$

and

$$\mathcal{D}_\mu \mathcal{D}_\nu - \mathcal{D}_\nu \mathcal{D}_\mu = 0.$$

Proof is by direct calculation•

Note that the volume form ℓ is constant w.r.t. these operators $\mathcal{D}_\mu$: $\mathcal{D}_\mu \ell = 0$, $\mu = 0, 1, 2, 3$.

14. Let differential forms $H \in \Lambda_1$ and $I, K \in \Lambda_2$ be such that

$$\begin{aligned} \mathcal{D}_\mu H &= 0, \quad \mathcal{D}_\mu I = 0, \quad \mathcal{D}_\mu K = 0, \quad \mu = 0, 1, 2, 3; \\ H^2 &= 1, \quad I^2 = K^2 = -1, \quad [H, I] = [H, K] = 0, \quad \{I, K\} = 0, \end{aligned} \quad (8)$$

where $\{I, K\} = IK + KI$.

We say that dx^μ are *primary generators* of Λ and H, ℓ, I, K are *secondary generators* of Λ .

The following differential forms from Λ are linear independent at every point $x \in \mathcal{V}$:

$$H, I, K, HI, HK, IK, HIK, \ell, \ell H, \ell I, \ell K, \ell HI, \ell HK, \ell IK, \ell HIK, 1. \quad (9)$$

These forms can be considered as basis forms of Λ .

In Minkowski space the concept of secondary generators of Clifford algebra $\mathcal{C}(1, 3)$ was presented in [3].

Theorem 2. *On the pseudo-Riemannian space $\mathcal{V}$ under consideration there exists the solution $H \in \Lambda_1$, $I, K \in \Lambda_2$, $B_\mu \in \Lambda_2 \top_1$ of the system of equation*

$$\mathcal{D}_\mu B_\nu - \mathcal{D}_\nu B_\mu + [B_\mu, B_\nu] = \frac{1}{2} C_{\mu\nu}, \quad (10)$$

$$\mathcal{D}_\mu H = 0, \quad \mathcal{D}_\mu I = \mathcal{D}_\mu K = 0, \quad (11)$$

$$H^2 = 1, \quad I^2 = K^2 = -1, \quad [H, I] = [H, K] = 0, \quad \{I, K\} = 0. \quad (12)$$

Proof. Consider local coordinates x^μ such that

$$g^{11} > 0, \quad \det \begin{vmatrix} g^{11} & g^{12} \\ g^{12} & g^{22} \end{vmatrix} < 0, \quad \det \begin{vmatrix} g^{11} & g^{12} & g^{13} \\ g^{12} & g^{22} & g^{23} \\ g^{13} & g^{23} & g^{33} \end{vmatrix} > 0. \quad (13)$$

Let us take

$$H = dx^1 / \sqrt{g^{11}},$$

$$I = \frac{-(-dx^2 \wedge dx^3 g^{11} - dx^1 \wedge dx^3 g^{12} + dx^1 \wedge dx^2 g^{13})}{\sqrt{g^{11}} \sqrt{-((g^{13})^2 g^{22}) + 2 g^{12} g^{13} g^{23} - g^{11} (g^{23})^2 - (g^{12})^2 g^{33} + g^{11} g^{22} g^{33}}} \quad (14)$$

$$K = -((dx^3 \wedge dx^4 (g^{12})^2 - dx^2 \wedge dx^4 g^{12} g^{13} + dx^2 \wedge dx^3 g^{12} g^{14} - dx^3 \wedge dx^4 g^{11} g^{22} + dx^1 \wedge dx^4 g^{13} g^{22} - dx^1 \wedge dx^3 g^{14} g^{22} + dx^2 \wedge dx^4 g^{11} g^{23} - dx^1 \wedge dx^4 g^{12} g^{23} + dx^1 \wedge dx^2 g^{14} g^{23} - dx^2 \wedge dx^3 g^{11} g^{24} + dx^1 \wedge dx^3 g^{12} g^{24} - dx^1 \wedge dx^2 g^{13} g^{24})(\sqrt{-g}))/\sqrt{(g^{12})^2 - g^{11} g^{22}}$$

which satisfy the equalities (12). Then we may consider the equalities (11) as the linear system of equations for the components $b_{\alpha\beta\mu} = b_{[\alpha\beta]\mu}$ of $B_\mu = \frac{1}{2}b_{\alpha\beta\mu}dx^\alpha \wedge dx^\beta$. The number of linear independent equations in this system of equations is equal to 24 and equal to the number of components $b_{\alpha\beta\mu}$, $\alpha < \beta$. A unique solution of this system of equations is written in Addendum. It can be checked that this solution also satisfies (10). This completes the proof.

Let us emphasize that $H = h_\mu dx^\mu$ has the form $H = dx^1/\sqrt{g^{11}}$ only in the fixed coordinates x^μ . We suppose that under a change of coordinates the set of values h_μ transforms as covector, i.e., $H \in \Lambda_1$. This remark is also true w.r.t. values $I, K \in \Lambda_2$, $B_\mu \in \Lambda_2 \mathbb{T}_1$ from (14) and from Addendum.

Note that if $\{H, I, K, B_\mu\}$ is a solution of the equations (10-12) and $S \in \text{Spin}(\mathcal{V})$, then $\{S^*HS, S^*IS, S^*KS, S^*B_\mu S - S^*\Upsilon_\mu S\}$ is also the solution of (10-12) (see Theorem 6).

15. Let us define the operation of Hermitian conjugation of differential forms

$$U^\dagger = HU^*H, \quad U \in \Lambda^\mathcal{K}.$$

Evidently,

$$U^{\dagger\dagger} = U, \quad (UV)^\dagger = V^\dagger U^\dagger$$

and $\alpha^\dagger = \bar{\alpha}$ for $\alpha \in \Lambda_0^\mathcal{K}$, where $\bar{\alpha}$ is the complex conjugated scalar function.

It is shown in ?? that in Minkowski space the operation of Hermitian conjugation of exterior forms closely connected to the operation of Hermitian conjugation of matrices.

16. Now we may define the operation $(\cdot, \cdot) : \Lambda^\mathcal{K} \times \Lambda^\mathcal{K} \rightarrow \Lambda_0^\mathcal{K}$ by the formula

$$(U, V) = \text{Tr}(UV^\dagger), \quad U, V \in \Lambda^\mathcal{K}.$$

This operation has all the properties of Hermitian scalar product at every point $x \in \mathcal{V}$

$$\begin{aligned} (\alpha U, V) &= \alpha(U, V), \quad (U, \alpha V) = \bar{\alpha}(U, V), \quad (U, V) = \overline{(V, U)}, \\ (U + W, V) &= (U, V) + (W, V), \quad (U, U) > 0 \text{ for } U \neq 0, \end{aligned}$$

where $U, V, W \in \Lambda^\mathcal{K}$, $\alpha \in \Lambda_0^\mathcal{K}$ and the bar means complex conjugation.

17. Let us summarize properties of the operators $\mathcal{D}_\mu$

- $\mathcal{D}_\mu \mathcal{D}_\nu - \mathcal{D}_\nu \mathcal{D}_\mu = 0$;
- $\mathcal{D}_\mu(UV) = (\mathcal{D}_\mu U)V + U\mathcal{D}_\mu V$;
- $\mathcal{D}_\mu(U + V) = \mathcal{D}_\mu U + \mathcal{D}_\mu V$;
- $\mathcal{D}_\mu \ell = 0$;
- $\mathcal{D}_\mu H = 0, \mathcal{D}_\mu I = 0, \mathcal{D}_\mu K = 0$;
- $\mathcal{D}_\mu(U^*) = (\mathcal{D}_\mu U)^*$;
- $\mathcal{D}_\mu(U^\dagger) = (\mathcal{D}_\mu U)^\dagger$;
- $\mathcal{D}_\mu(\star U) = \star(\mathcal{D}_\mu U)$;
- $\mathcal{D}_\mu(\text{Tr} U) = \text{Tr}(\mathcal{D}_\mu U)$;
- $\mathcal{D}_\mu(U, V) = (\mathcal{D}_\mu U, V) + (U, \mathcal{D}_\mu V)$.

18. In what follows we use the four spaces of differential forms $\Lambda^\mathcal{C}, \Lambda, \Lambda_{\text{ev}}^\mathcal{C}, \Lambda_{\text{ev}}$. Each of these spaces can be considered at every point $x \in \mathcal{V}$ as a bialgebra with two products, namely the exterior product and Clifford product. We use notation Ω for any of these spaces. The operation $(\cdot, \cdot)$ converts a space Ω into the unitary space at every point $x \in \mathcal{V}$.

19. We say that the differential form $U \in \Omega$ is *Hermitian* if $U^\dagger = U$ and *anti-Hermitian* if $U^\dagger = -U$. Every differential form $U \in \Omega$ can be decomposed into Hermitian and anti-Hermitian parts

$$U = \frac{1}{2}(U + U^\dagger) + \frac{1}{2}(U - U^\dagger).$$

Let us denote

$$\Omega_- = \{U \in \Omega : U^\dagger = -U\}.$$

This set of differential forms is closed w.r.t. commutator $[\cdot, \cdot]$ and can be considered as a real Lie algebra.

20. Let us introduce two differential forms N, E

$$\begin{aligned} N &= i & \text{for } \Omega = \Lambda^c, \Lambda_{\text{ev}}^c; \\ N &= \ell & \text{for } \Omega = \Lambda; \\ N &= I & \text{for } \Omega = \Lambda_{\text{ev}}; \\ E &= 1 & \text{for } \Omega = \Lambda^c, \Lambda; \\ E &= H & \text{for } \Omega = \Lambda_{\text{ev}}^c, \Lambda_{\text{ev}} \end{aligned} \tag{15}$$

with the following properties:

$$\begin{aligned} \mathcal{D}_\mu N &= 0, \quad \mathcal{D}_\mu E = 0, \quad \mu = 0, 1, 2, 3; \\ N^2 &= -1, \quad E^2 = 1, \quad [N, E] = 0; \\ N^\dagger &= -N, \quad E^\dagger = E. \end{aligned} \tag{16}$$

21. Let us define the set of differential forms

$$L_{\max} = \{U \in \Omega_- : [U, N] = [U, E] = 0\}.$$

This set of differential forms is closed w.r.t. the commutator $[\cdot, \cdot]$ and can be considered as Lie algebra.

Let $t_1, \dots, t_p$ be generators of $L_{\max}$ such that

$$\begin{aligned} \mathcal{D}_\mu t_k &= 0, \quad t_k^\dagger = -t_k, \quad (t_k, t_l) = \delta_{kl}, \\ [t_k, N] &= [t_k, E] = 0, \quad [t_k, t_l] = \sum_{q=1}^p c_{kl}^q t_q \quad k, l = 1, \dots, p, \end{aligned} \tag{17}$$

where c_{kl}^q are real structure constants of the Lie algebra $L_{\max}$. Hence

$$L_{\max} = L(t_1, \dots, t_p) = \left\{ \sum_{k=1}^p f_k t_k \right\},$$

where $f_k = f_k(x)$, $x \in \mathcal{V}$ are smooth real valued functions from $C^\infty(\mathcal{V}, \mathcal{R})$.

22. Let L_0 be a subalgebra of the Lie algebra L_{max} such that the first d ($d \leq p$) generators $t_1, \dots, t_d$ of L_{max} are the generators of L_0

$$L_0 = L(t_1, \dots, t_d) = \left\{ \sum_{k=1}^n f_k t_k \right\}$$

and

$$\begin{aligned} [t_k, t_l] &= \sum_{q=1}^d c_{kl}^q t_q, \quad k, l = 1, \dots, d, \\ [t_k, t_r] &= \sum_{q=d+1}^p c_{kr}^q t_q, \quad k = 1, \dots, d; \quad r = d+1, \dots, p. \end{aligned}$$

23. Let us define the set of differential forms

$$G_0 = \exp L_0 = \{ \exp u : u \in L_0 \},$$

where

$$\exp u = 1 + \sum_{k=1}^{\infty} \frac{1}{k!} u^k.$$

G_0 is a unitary ($U^{-1} = U^\dagger$) Lie group w.r.t. Clifford product and L_0 is the real Lie algebra of the Lie group G_0 .

24. Let us take

$$\alpha^\mu = H dx^\mu \in \Lambda_{\text{ev}} \mathbb{T}^1$$

such that $(\alpha^\mu)^\dagger = \alpha^\mu$.

2 Lie algebras of anti-Hermitian differential forms.

Let $\Omega = \Lambda^{\mathcal{C}}$. Consider the following 16 differential forms from Ω :

$$iH, I, K, HI, HK, IK, HIK, \ell, i\ell H, i\ell I, i\ell K, \ell HI, \ell HK, i\ell IK, \ell HIK, i \quad (18)$$

and denote them by $T_1, \dots, T_{16}$. It is not hard to check that

$$\mathcal{D}_\mu T_k = 0, \quad T_k^\dagger = -T_k, \quad (T_k, T_l) = \delta_{kl}, \quad [T_k, T_l] = c_{kl}^q T_q, \quad (19)$$

where c_{kl}^q are real structure constants.

Every element $U \in \Omega_-$ can be written in the form

$$U = \sum_{k=1}^{16} (U, T_k) T_k.$$

Hence these differential forms T_k are generators of the real Lie algebra Ω_-

$$\Omega_- = L(T_1, \dots, T_{16}) = \left\{ \sum_{k=1}^{16} f_k T_k \right\},$$

where $f_k = f_k(x)$, $x \in \mathcal{V}$ are real valued functions from $C^\infty(\mathcal{V}, \mathcal{R})$; It can be shown [1] that the Lie algebra Ω_- (considered at every point $x \in \mathcal{V}$) is isomorphic to the Lie algebra $u(4) \simeq u(1) \oplus su(4)$ of anti-Hermitian 4×4 -matrices. So

$$\Omega_- = L(T_{16}) \oplus L(T_1, \dots, T_{15}),$$

where $L(T_{16})$ is isomorphic to $u(1)$ and $L(T_1, \dots, T_{15})$ is isomorphic to $su(4)$ (at every point $x \in \mathcal{V}$).

For $\Omega = \Lambda^c$ we have $N = i$, $E = 1$. Consequently, $L_{max} = \Omega_-$. Continuing this line of reasoning, we find real Lie algebras Ω_- and L_{max} for the cases $\Omega = \Lambda, \Lambda_{ev}^c, \Lambda_{ev}$.

Let us summarize our consideration of real Lie algebras of differential forms.

- If $\Omega = \Lambda^c$, then $N = i$, $E = 1$ and

$$\Omega_- = L_{max} = L(i) \oplus L(iH, I, K, HI, \dots, \ell HIK) \simeq u(1) \oplus su(4).$$

- If $\Omega = \Lambda$, then $N = \ell$, $E = 1$ and

$$\begin{aligned} \Omega_- &= L(I, K, HI, HK, IK, HIK, \ell, \ell HI, \ell HK, \ell HIK) \simeq sp(2), \\ L_{max} &= L(\ell) \oplus L(I, K, IK) \simeq u(1) \oplus su(2), \end{aligned}$$

where $sp(2)$ is the Lie algebra of the symplectic unitary Lie group.

- If $\Omega = \Lambda_{ev}^c$, then $N = i$, $E = H$ and

$$\begin{aligned} \Omega_- &= L(i) \oplus L(\ell) \oplus L(I, K, IK, i\ell I, i\ell K, i\ell IK) \\ &\simeq u(1) \oplus u(1) \oplus su(2) \oplus su(2), \\ L_{max} &= L(i) \oplus L(I, K, IK) \simeq u(1) \oplus su(2), \end{aligned}$$

- If $\Omega = \Lambda_{\text{ev}}$, then $N = I$, $E = H$ and

$$\begin{aligned}\Omega_- &= L(I) \oplus L(\ell) \simeq u(1) \oplus u(1), \\ L_{\max} &= L(I) \simeq u(1),\end{aligned}$$

Let us consider differential forms $t_1, \dots, t_{16}$ such that $t_k = \sum_{p=1}^{16} r_k^p T_p$, where real constants r_k^p are elements of an orthogonal 16×16 -matrix. Evidently

$$\mathcal{D}_\mu t_k = 0, \quad t_k^\dagger = -t_k, \quad (t_k, t_l) = \delta_{kl}$$

and the set of differential forms $\{t_1, \dots, t_{16}\}$ is another set of generators of the real Lie algebra $\Omega_- \simeq u(4)$ in case $\Omega = \Lambda^{\mathcal{C}}$. We claim that this construction allow us to describe a Lie subalgebra of Ω_- that is isomorphic to the Lie algebra $su(3)$. Indeed, let us take the following generators of Ω_- :

$$\begin{aligned}t_1 &= (HIK + IK)/\sqrt{2}, \\ t_2 &= (HK + K)/\sqrt{2}, \\ t_3 &= (-HI - I)/\sqrt{2}, \\ t_4 &= (\ell - \ell Ii)/\sqrt{2}, \\ t_5 &= (\ell HI + \ell Hi)/\sqrt{2}, \\ t_6 &= (-\ell HK + \ell IKi)/\sqrt{2}, \\ t_7 &= (-\ell HIK - \ell Ki)/\sqrt{2}, \\ t_8 &= (-HI - 2Hi + I)/\sqrt{6}, \\ t_9 &= (\ell HK + \ell IKi)/\sqrt{2}, \\ t_{10} &= (-\ell HIK + \ell Ki)/\sqrt{2}, \\ t_{11} &= (\ell + \ell Ii)/\sqrt{2}, \\ t_{12} &= (-\ell HI + \ell Hi)/\sqrt{2}, \\ t_{13} &= (-HIK + IK)/\sqrt{2}, \\ t_{14} &= (-HK + K)/\sqrt{2}, \\ t_{15} &= (HI - Hi - I)/\sqrt{3}, \\ t_{16} &= i,\end{aligned} \tag{20}$$

which are expressed via generators $T_1, \dots, T_{16}$ from (18) with the aid of the orthogonal matrix. It can be easily checked that differential forms $t_1, \dots, t_8$

are the generators of the real Lie algebra $L_0 = L(t_1, \dots, t_8)$ isomorphic to the Lie algebra $su(3)$ and

$$\begin{aligned} [t_k, t_l] &= \sum_{q=1}^8 c_{kl}^q t_q, \quad k, l = 1, \dots, 8, \\ [t_k, t_r] &= \sum_{q=9}^{15} c_{kr}^q t_q, \quad k = 1, \dots, 8; \quad r = 9, \dots, 16. \end{aligned}$$

3 Main equations.

Now we may write down the main system of equations

$$\mathcal{D}_\mu B_\nu - \mathcal{D}_\nu B_\mu + [B_\mu, B_\nu] = \frac{1}{2} C_{\mu\nu}, \quad (21)$$

$$\mathcal{D}_\mu N = 0, \quad \mathcal{D}_\mu E = 0, \quad \mu = 0, 1, 2, 3; \quad (22)$$

$$N^2 = -1, \quad E^2 = 1, \quad [N, E] = 0; \quad N^\dagger = -N, \quad E^\dagger = E. \quad (23)$$

$$dx^\mu (\mathcal{D}_\mu \Psi + \Psi A_\mu + B_\mu \Psi) N - m \Psi E = 0, \quad (24)$$

$$\mathcal{D}_\mu A_\nu - \mathcal{D}_\nu A_\mu - [A_\mu, A_\nu] = F_{\mu\nu}, \quad (25)$$

$$\frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} F^{\mu\nu}) - [A_\mu, F^{\mu\nu}] = J^\nu, \quad (26)$$

$$J^\nu = \frac{1}{4} \sum_{k=1}^d (E \{E, \{N, \Psi^\dagger \alpha^\nu \Psi\}\}, t_k) t_k, \quad (27)$$

where $\Psi \in \Omega$, $A_\mu \in L_0 \top_1$, $F_{\mu\nu} \in L_0 \top_2$, $F^{\mu\nu} = g^{\mu\alpha} g^{\nu\beta} F_{\alpha\beta} \in L_0 \top_2$, $B_\mu \in \Lambda_2 \top_1$, m is a real constant. The values $J^\nu = J^\nu(\Psi, E, N) \in L_0 \top^1$ also can be defined with the aid of the following equalities:

$$\begin{aligned} J_{(1)}^\nu &= \Psi^\dagger \alpha^\nu \Psi \in \Omega \top^1 : (J_{(1)}^\nu)^\dagger = J_{(1)}^\nu. \\ J_{(2)}^\nu &= \frac{1}{2} \{N, J_{(1)}^\nu\} \in \Omega_- \top^1 : [J_{(2)}^\nu, N] = 0, \end{aligned} \quad (28)$$

$$\begin{aligned}
J_{(3)}^\nu &= \frac{1}{2}E\{E, J_{(2)}^\nu\} \in L_{max} \top^1 : [J_{(3)}^\nu, N] = [J_{(3)}^\nu, E] = 0, \\
J^\nu &= \sum_{k=1}^d (J_{(3)}^\nu, t_k) t_k \in L_0 \top^1,
\end{aligned}$$

where $t_1, \dots, t_d$ are the generators (17) of the real Lie algebra L_0 (see subsection 22 of section 1).

We say that the equation (24) is a *Dirac type tensor equation* and the equations (25,26) are *Yang-Mills equations*.

The full set of $\{A_\mu, F_{\mu\nu}\}$ is called *the Yang-Mills field*; A_μ is potential and $F_{\mu\nu}$ is strength of the Yang-Mills field.

In the Standard Model of elementary particles three real Lie algebras are of special interest. Namely the Lie algebra $su(3)$ is used in Quantum Chromodynamics, $u(1) \oplus su(2)$ is used in Electroweak Theory, and $u(1)$ is used in Quantum Electrodynamics. Taking these Lie algebras into account, we consider five special cases of equations (21-27).

(i) $\Omega = \Lambda^c, N = i, E = 1,$

$$L_0 = L_{max} \simeq u(1) \oplus su(4), \quad J^\nu = i\Psi^\dagger \alpha^\nu \Psi;$$

(ii) $\Omega = \Lambda^c, N = i, E = 1,$

$$L_0 = L(t_1, \dots, t_8) \simeq su(3), \quad J^\nu = \sum_{k=1}^8 (i\Psi \alpha^\nu \Psi, t_k) t_k,$$

where $t_1, \dots, t_8$ from (20);

(iii) $\Omega = \Lambda_{ev}^c, N = i, E = H,$

$$L_0 = L_{max} \simeq u(1) \oplus su(2), \quad J^\nu = \frac{1}{2}H\{H, i\Psi^\dagger \alpha^\nu \Psi\};$$

(iv) $\Omega = \Lambda, N = \ell, E = 1,$

$$L_0 = L_{max} \simeq u(1) \oplus su(2), \quad J^\nu = \frac{1}{2}\{\ell, \Psi^\dagger \alpha^\nu \Psi\};$$

$$(v) \Omega = \Lambda_{ev}, N = I, E = H,$$

$$L_0 = L_{max} = L(I) \simeq u(1), \quad J^\nu = \frac{1}{4}H\{H, \{I, \Psi^\dagger \alpha^\nu \Psi\}\}.$$

The full set of $\{\Psi, E, N\}$ is called *the wave field* of the equations (22-24).

In the system of equations (21-27) we consider the wave field $\{\Psi, E, N\}$, the Yang-Mills field $\{A_\mu, F_{\mu\nu}\}$, and the field B_μ as unknown values.

4 Nonabelian charge conservation laws.

Theorem 3. *If the values $\Psi \in \Omega$, $A_\mu \in L_0\mathbb{T}_1$, $B_\mu \in \Lambda_2\mathbb{T}_1$, $N \in \Omega$, $E \in \Omega$ satisfy Dirac type tensor equation (24) and the values $J^\mu \in L_0\mathbb{T}^1$ are defined in (27) or, equivalently, in (28), then J^μ satisfy the equality*

$$\frac{1}{\sqrt{-g}}\mathcal{D}_\mu(\sqrt{-g}J^\mu) - [A_\mu, J^\mu] = 0. \quad (29)$$

The equality (29) is called *the nonabelian charge conservation law* for the Dirac type tensor equation.

Proof. Let us multiply the equation (24) from the left by H and from the right by $-N$. We denote the left hand side of resulting equation by

$$Q = \alpha^\mu(\mathcal{D}_\mu\Psi + \Psi A_\mu + B_\mu\Psi) + mH\Psi EN. \quad (30)$$

Then

$$Q^\dagger = (\mathcal{D}_\mu\Psi^\dagger - A_\mu\Psi^\dagger + \Psi^\dagger B_\mu^\dagger)\alpha^\mu - mNE\Psi^\dagger H.$$

Consider the expression

$$\begin{aligned} Y_{(1)} &= \Psi^\dagger Q + Q^\dagger \Psi \\ &= \mathcal{D}_\mu(\Psi^\dagger \alpha^\mu \Psi) + \Psi^\dagger(-\mathcal{D}_\mu \alpha^\mu + \alpha^\mu B_\mu + B_\mu^\dagger \alpha^\mu)\Psi - [A_\mu, \Psi^\dagger \alpha^\mu \Psi] + m[\Psi^\dagger H\Psi, EN] \\ &= \mathcal{D}_\mu(\Psi^\dagger \alpha^\mu \Psi) + \Gamma_{\mu\nu}^\mu \Psi^\dagger \alpha^\nu \Psi - [A_\mu, \Psi^\dagger \alpha^\mu \Psi] + m[\Psi^\dagger H\Psi, EN] \\ &= \frac{1}{\sqrt{-g}}\mathcal{D}_\mu(\sqrt{-g}J_{(1)}^\mu) - [A_\mu, J_{(1)}^\mu] + m[\Psi^\dagger H\Psi, EN]. \end{aligned}$$

Here we use the formulas

$$\begin{aligned}\mathcal{D}_\mu \alpha^\mu &= -\Gamma_{\mu\nu}^\mu \alpha^\nu + \alpha^\mu B_\mu + B_\mu^\dagger \alpha^\mu, \\ \mathcal{D}_\mu J_{(1)}^\mu + \Gamma_{\mu\nu}^\mu J_{(1)}^\nu &= \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} J_{(1)}^\mu),\end{aligned}$$

see [2].

If we take

$$\begin{aligned}Y_{(2)} &= \frac{1}{2} \{N, Y_{(1)}\}, \\ Y_{(3)} &= \frac{1}{2} E \{E, Y_{(2)}\},\end{aligned}$$

then we get

$$Y_{(3)} = \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} J_{(3)}^\mu) - [A_\mu, J_{(3)}^\mu] + \frac{1}{4} m E \{E, \{N, [\Psi^\dagger H \Psi, EN]\}\}.$$

Using the equalities $N^2 = -1$, $E^2 = 1$, $[N, E] = 0$, it is easy to check that

$$E \{E, \{N, [\Psi^\dagger H \Psi, EN]\}\} \equiv 0.$$

Hence,

$$Y_{(3)} = \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} J_{(3)}^\mu) - [A_\mu, J_{(3)}^\mu].$$

Finally, we take

$$Y = \sum_{k=1}^d (Y_{(3)}, t_k) t_k,$$

where the generators $t_1, \dots, t_d$ of the Lie algebra L_0 are defined in the subsection 22 of section 1. In this case we obtain

$$Y = \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} J^\mu) - [A_\mu, J^\mu].$$

By assumption, Ψ, N, E, A_μ, B_μ satisfy (24). Consequently, $Q = Y_{(1)} = Y_{(2)} = Y_{(3)} = Y = 0$ and J^μ satisfy equality (29). This completes the proof.

Theorem 4. *Let us denote the left hand side of the equation (26) by R^ν*

$$R^\nu = \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} F^{\mu\nu}) - [A_\mu, F^{\mu\nu}],$$

where $F_{\mu\nu}$ satisfy (25). Then

$$\frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} R^\mu) - [A_\mu, R^\mu] = 0.$$

The proof is by direct calculation.

This theorem means that the equation (26) is consistent with the non-abelian charge conservation law (29).

5 Unitary and Spin gauge symmetries.

Theorem 5. *Suppose that the following conditions hold:*

1. *The values*

$$\begin{aligned} \Psi &\in \Omega, A_\mu \in L_0 \mathbb{T}_1, F_{\mu\nu} \in L_0 \mathbb{T}_2, J^\nu \in L_0 \mathbb{T}^1, \\ E &\in \Omega, N \in \Omega, B_\mu \in \Lambda_2 \mathbb{T}_1, C_{\mu\nu} \in \Lambda_2 \mathbb{T}_2 \end{aligned} \quad (31)$$

and operators $\mathcal{D}_\mu = \Upsilon_\mu - [B_\mu, \cdot]$ satisfy equations (21-27).

2. $U = U(x)$ is a smooth map $\mathcal{V} \rightarrow G_0$, where the Lie group G_0 is defined in subsection 23 of section 1.

Then the values with acute

$$\begin{aligned} \acute{\Psi} &= \Psi U, \acute{A}_\mu = U^{-1} A_\mu U - U^{-1} \mathcal{D}_\mu U, \acute{F}_{\mu\nu} = U^{-1} F_{\mu\nu} U, \acute{J}^\nu = U^{-1} J^\nu U, \\ \{\acute{E}, \acute{N}, \acute{B}_\mu, \acute{C}_{\mu\nu}\} &= \{E, N, B_\mu, C_{\mu\nu}\} \end{aligned} \quad (32)$$

and the operators $\acute{\mathcal{D}}_\mu = \mathcal{D}_\mu$ satisfy the same equations (21-27).

The proof is straightforward.

Theorem 6. *If the values (31) and operators $\mathcal{D}_\mu$ satisfy the equations (21-27) and $S = S(x)$ is a smooth map $\mathcal{V} \rightarrow \text{Spin}(\mathcal{V})$, then the values with check*

$$\begin{aligned}\check{\Psi} &= \Psi S, \check{A}_\mu = S^{-1} A_\mu S, \check{F}_{\mu\nu} = S^{-1} F_{\mu\nu} S, \check{J}^\nu = S^{-1} J^\nu S, \\ \check{E} &= S^{-1} E S, \check{N} = S^{-1} N S, \check{B}_\mu = S^{-1} B_\mu S - S^{-1} \Upsilon_\mu S, \check{C}_{\mu\nu} = C_{\mu\nu}\end{aligned}\quad (33)$$

and the operators $\check{\mathcal{D}}_\mu = \Upsilon_\mu - [\check{B}_\mu, \cdot]$ satisfy the same equations (21-27).

The proof is by direct calculation. Note that for values with check the operation of Hermitian conjugation is defined by

$$\check{U}^\dagger = \check{H} \check{U}^* \check{H},$$

where $\check{H} = S^{-1} H S$.

Invariance of the equation (21) is proved in [2].

6 Lagrangians.

Denote

$$P = dx^\mu (\mathcal{D}_\mu \Psi + \Psi A_\mu + B_\mu \Psi) N - m \Psi E. \quad (34)$$

We have $\Psi \in \Omega$

$$\begin{aligned}\Psi &= \psi + \psi_\mu dx^\mu + \frac{1}{2!} \psi_{\mu_1 \mu_2} dx^{\mu_1} \wedge dx^{\mu_2} + \dots + \frac{1}{4!} \psi_{\mu_1 \dots \mu_4} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_4}, \\ \psi_{\mu_1 \dots \mu_k} &= \psi_{[\mu_1 \dots \mu_k]} = p_{\mu_1 \dots \mu_k} + i q_{\mu_1 \dots \mu_k},\end{aligned}$$

where $p_{\mu_1 \dots \mu_k}$, $q_{\mu_1 \dots \mu_k}$ are real valued components of a covariant antisymmetric tensor fields of rank k and

$$\begin{aligned}q_{\mu_1 \dots \mu_k} &\equiv 0 \quad (k = 0, \dots, 4) \quad \text{for} \quad \Omega = \Lambda, \Lambda_{\text{ev}}, \\ \psi_\mu, \psi_{\mu_1 \mu_2 \mu_3} &\equiv 0 \quad \text{for} \quad \Omega = \Lambda_{\text{ev}}^c, \Lambda_{\text{ev}}.\end{aligned}$$

Consider the Lagrangian (Lagrangian density)

$$\mathcal{L}_0 = \frac{\sqrt{-g}}{4} \text{Tr}(H(\Psi^* P + P^* \Psi)).$$

A Dirac type tensor equation can be derived from $\mathcal{L}_0$ with the aid of the following formulas:

$$\begin{aligned} u^{\mu_1 \dots \mu_k} &= \left(\frac{\partial \mathcal{L}_0}{\partial p_{\mu_1 \dots \mu_k}} - \partial_\nu \frac{\partial \mathcal{L}_0}{\partial p_{\mu_1 \dots \mu_k, \nu}} \right) + i \left(\frac{\partial \mathcal{L}_0}{\partial q_{\mu_1 \dots \mu_k}} - \partial_\nu \frac{\partial \mathcal{L}_0}{\partial q_{\mu_1 \dots \mu_k, \nu}} \right), \\ u_{\nu_1 \dots \nu_k} &= u_{[\nu_1 \dots \nu_k]} = g_{\mu_1 \nu_1} \dots g_{\mu_k \nu_k} u^{\mu_1 \dots \mu_k}, \\ U &= \sum_{k=0}^4 \frac{1}{k!} u_{\nu_1 \dots \nu_k} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_k}, \\ \dot{P} &= \frac{1}{\sqrt{-g}} U H, \end{aligned}$$

where $p_{\mu_1 \dots \mu_k, \nu} = \partial_\nu p_{\mu_1 \dots \mu_k}$, $q_{\mu_1 \dots \mu_k, \nu} = \partial_\nu q_{\mu_1 \dots \mu_k}$. Now we may check that $\dot{P} \equiv P$.

Let us consider the Lagrangian for Yang-Mills equations. Denote

$$\begin{aligned} F_{\mu\nu} &= \mathcal{D}_\mu A_\nu - \mathcal{D}_\nu A_\mu - [A_\mu, A_\nu], \\ F^{\alpha\beta} &= g^{\alpha\mu} g^{\beta\nu} F_{\mu\nu}, \\ Y^\nu &= \frac{1}{\sqrt{-g}} \mathcal{D}_\mu (\sqrt{-g} F^{\mu\nu}) - [A_\mu, F^{\mu\nu}] - J^\nu, \end{aligned}$$

where J^ν are from (27) and $A_\mu = a_\mu^k t_k$, ($t_k = t^k$ are defined in the subsection 22 of section 1).

The Yang-Mills Lagrangian has the form

$$\mathcal{L}_1 = \frac{\sqrt{-g}}{4} \text{Tr}(F_{\mu\nu} F^{\mu\nu})$$

and the complete Lagrangian for the Dirac type tensor equation together with Yang-Mills equations

$$\mathcal{L} = \mathcal{L}_1 + 2n\mathcal{L}_0.$$

Here $n = 4$ is the dimension of space-time. Denote

$$\begin{aligned} y_k^\mu &= \frac{\partial \mathcal{L}}{\partial a_\mu^k} - \partial_\nu \frac{\partial \mathcal{L}}{\partial a_{\mu, \nu}^k}, \\ \dot{Y}^\nu &= \frac{1}{\sqrt{-g}} y_k^\mu t^k. \end{aligned}$$

Now we may check that

$$\dot{Y}^\nu \equiv Y^\nu.$$

Hence the Dirac type tensor equation and the Yang-Mills equations can be derived from the Lagrangian $\mathcal{L}$.

We hope that equations (21-27) can be used to describe fermions in space-time with presence of gravitation. From this point of view it is reasonable to add the Einstein-Hilbert Lagrangian $\sqrt{-g} R$ to $\mathcal{L}$ and variate the resulting Lagrangian w.r.t. components of metric tensor $g_{\mu\nu}$.

7 A geometrical interpretation of results.

We consider a pseudo-Riemannian space $\mathcal{V}$ with the Levi-Civita connection $\Gamma^\lambda_{\mu\nu}$, with the covariant derivatives ∇_μ , with the Upsilon derivatives Υ_μ , and with the curvature tensor $R_{\alpha\beta\mu\nu}$ defined in section 1. Suppose that a new structure on $\mathcal{V}$ is given. Namely the affine connection $\check{\Gamma}^\lambda_{\mu\nu}$. We get definitions of the covariant derivatives $\check{\nabla}_\mu$, the Upsilon derivatives $\check{\Upsilon}_\mu$, and the curvature tensor $\check{R}_{\alpha\beta\mu\nu}$, replacing $\Gamma^\lambda_{\mu\nu}$ by $\check{\Gamma}^\lambda_{\mu\nu}$ in the corresponding definitions in section 1. We suppose that the affine connection $\check{\Gamma}^\lambda_{\mu\nu}$ is metric compatible

$$\check{\nabla}_\kappa g_{\mu\nu} = 0, \quad \check{\nabla}_\kappa g^{\mu\nu} = 0.$$

It is convenient to introduce the tensor

$$K^\lambda_{\mu\nu} = \check{\Gamma}^\lambda_{\mu\nu} - \Gamma^\lambda_{\mu\nu},$$

which we, following [10], will call contorsion. It is easy to see that the affine connection is metric compatible iff $K_{\nu\mu\lambda} = -K_{\lambda\mu\nu}$. Torsion is expressed via contorsion as

$$T^\lambda_{\mu\nu} = K^\lambda_{\mu\nu} - K^\lambda_{\nu\mu}.$$

Conversely, the contorsion of a metric compatible connection is expressed via torsion as

$$K^\lambda_{\mu\nu} = \frac{1}{2}(T^\lambda_{\mu\nu} + T^\lambda_{\nu\mu} + T^\lambda_{\mu\lambda})$$

(see [10], formula (7.35)).

So we arrive at the affine space $\{\mathcal{M}, g_{\mu\nu}, K^\lambda{}_{\mu\nu}\}$. Let us define the tensors

$$\begin{aligned} b_{\alpha\beta\mu} &= -\frac{1}{2}K_{\alpha\mu\beta}, \\ B_\mu &= \frac{1}{2}b_{\alpha\beta\mu}dx^\alpha \wedge dx^\beta \in \Lambda^2\mathbb{T}_1, \\ C_{\mu\nu} &= \frac{1}{2}R_{\alpha\beta\mu\nu}dx^\alpha \wedge dx^\beta \in \Lambda^2\mathbb{T}_2. \end{aligned}$$

Theorem 7. *If $U \in \Lambda$, then*

$$\check{\Upsilon}_\mu U = \Upsilon_\mu U - [B_\mu, U].$$

Proof follows from the formula

$$K^\nu{}_{\mu\lambda}dx^\lambda = [B_\mu, dx^\nu],$$

which can be easily checked.

Theorem 8. *(F.E.Burstall, A.D.King, N.G.Marchuk, D.G.Vassiliev) The following equality holds*

$$\Upsilon_\mu B_\nu - \Upsilon_\nu B_\mu - [B_\mu, B_\nu] = \frac{1}{2}C_{\mu\nu}$$

iff

$$\check{R}_{\alpha\beta\mu\nu} = 0.$$

Proof. Suppose that the tensors $q_{\alpha\beta\mu\nu}$ and $\check{R}_{\alpha\beta\mu\nu}$ are such that

$$\frac{1}{2}q_{\alpha\beta\mu\nu}dx^\alpha \wedge dx^\beta = \Upsilon_\mu B_\nu - \Upsilon_\nu B_\mu - [B_\mu, B_\nu] - \frac{1}{2}C_{\mu\nu}$$

and

$$\check{R}_{\alpha\lambda\mu\nu} = g_{\kappa\alpha}(\partial_\mu \check{\Gamma}^\kappa{}_{\nu\lambda} - \partial_\nu \check{\Gamma}^\kappa{}_{\mu\lambda} + \check{\Gamma}^\kappa{}_{\mu\eta} \check{\Gamma}^\eta{}_{\nu\lambda} - \check{\Gamma}^\kappa{}_{\nu\eta} \check{\Gamma}^\eta{}_{\mu\lambda}). \quad (35)$$

Then

$$\check{R}_{\alpha\beta\mu\nu} = -2q_{\alpha\beta\mu\nu}.$$

This completes the proof.

8 Space dimensions $n = 2, 3$.

Let $\mathcal{M}_3$ be a three dimensional differentiable manifolds with local coordinates x^1, x^2, x^3 and with the smooth metric tensor $g_{\mu\nu} = g_{\nu\mu}$, $(\mu, \nu = 1, 2, 3)$ that satisfy (13). Denote by $\mathcal{V}_3$ the pseudo-Riemannian space $\{\mathcal{M}_3, g\}$ with the Levi-Civita connection $\Gamma^\lambda_{\mu\nu}$, with the covariant derivatives ∇_μ , with the Upsilon derivatives Υ_μ , and with the curvature tensor $R_{\alpha\beta\mu\nu}$.

Theorem 10. *On the pseudo-Riemannian space $\mathcal{V}_3$ there exists the solution $H \in \Lambda_1$, $I \in \Lambda_2$, $B_\mu \in \Lambda_2 \top_1$ of the system of equations that consists of (21) and*

$$\mathcal{D}_\mu H = 0, \quad \mathcal{D}_\mu I = 0, \quad (36)$$

$$H^2 = 1, \quad I^2 = -1, \quad [H, I] = 0, \quad (37)$$

where $\mathcal{D}_\mu = \Upsilon_\mu - [B_\mu, \cdot]$.

Proof. Evidently H, I from (14) satisfy (37). It can be checked that the following components $b_{\alpha\beta\mu} = b_{[\alpha\beta]\mu}$ of $B_\mu = \frac{1}{2}b_{\alpha\beta\mu}dx^\alpha \wedge dx^\beta$ satisfy (21) and (36):

$$b_{121} = (\partial_1 g_{33} g_{13} g_{22} g_{23} - \partial_1 g_{33} g_{12} g_{23}^2 - 2 \partial_1 g_{23} g_{13} g_{22} g_{33} + 2 \partial_1 g_{23} g_{12} g_{23} g_{33} + \partial_1 g_{22} g_{13} g_{23} g_{33} - \partial_2 g_{11} g_{23}^2 g_{33} - \partial_1 g_{22} g_{12} g_{33}^2 + \partial_2 g_{11} g_{22} g_{33}^2) / (4 g_{33} (-g_{23}^2 + g_{22} g_{33}))$$

$$b_{122} = (\partial_2 g_{33} g_{13} g_{22} g_{23} - \partial_2 g_{33} g_{12} g_{23}^2 - 2 \partial_2 g_{23} g_{13} g_{22} g_{33} + 2 \partial_2 g_{23} g_{12} g_{23} g_{33} + \partial_2 g_{22} g_{13} g_{23} g_{33} + \partial_1 g_{22} g_{23}^2 g_{33} - 2 \partial_2 g_{12} g_{23}^2 g_{33} - \partial_2 g_{22} g_{12} g_{33}^2 - \partial_1 g_{22} g_{22} g_{33}^2 + 2 \partial_2 g_{12} g_{22} g_{33}^2) / (4 g_{33} (-g_{23}^2 + g_{22} g_{33}))$$

$$b_{123} = (\partial_3 g_{33} g_{13} g_{22} g_{23} - \partial_3 g_{33} g_{12} g_{23}^2 - 2 \partial_3 g_{23} g_{13} g_{22} g_{33} + 2 \partial_3 g_{23} g_{12} g_{23} g_{33} + \partial_3 g_{22} g_{13} g_{23} g_{33} + \partial_1 g_{23} g_{23}^2 g_{33} - \partial_2 g_{13} g_{23}^2 g_{33} - \partial_3 g_{12} g_{23}^2 g_{33} - \partial_3 g_{22} g_{12} g_{33}^2 - \partial_1 g_{23} g_{22} g_{33}^2 + \partial_2 g_{13} g_{22} g_{33}^2 + \partial_3 g_{12} g_{22} g_{33}^2) / (4 g_{33} (-g_{23}^2 + g_{22} g_{33}))$$

$$b_{131} = (-\partial_1 g_{33} g_{13} + \partial_3 g_{11} g_{33}) / (4 g_{33})$$

$$b_{132} = (-\partial_2 g_{33} g_{13} - \partial_1 g_{23} g_{33} + \partial_2 g_{13} g_{33} + \partial_3 g_{12} g_{33}) / (4 g_{33})$$

$$b_{133} = (-\partial_3 g_{33} g_{13} - \partial_1 g_{33} g_{33} + 2 \partial_3 g_{13} g_{33}) / (4 g_{33})$$

$$b_{231} = (-\partial_1 g_{33} g_{23} - \partial_1 g_{23} g_{33} + \partial_2 g_{13} g_{33} - \partial_3 g_{12} g_{33}) / (4 g_{33})$$

$$b_{232} = (-\partial_2 g_{33} g_{23} - \partial_3 g_{22} g_{33}) / (4 g_{33})$$

$$b_{233} = (-\partial_3 g_{33} g_{23} + \partial_2 g_{33} g_{33} - 2 \partial_3 g_{23} g_{33}) / (4 g_{33})$$

Note that $\{S^* H S, S^* I S, S^* B_\mu S - S^* \Upsilon_\mu S\}$ also satisfy (36-37), where $S \in \text{Spin}(\mathcal{V}_3)$.

Let $\mathcal{M}_2$ be a two dimensional differentiable manifolds with local coordinates x^1, x^2 and with the smooth metric tensor $g_{\mu\nu} = g_{\nu\mu}$, $(\mu, \nu = 1, 2)$ such that $g_{11} > 0$, $g_{11}g_{22} - (g_{12})^2 < 0$. Denote by $\mathcal{V}_2$ the pseudo-Riemannian space $\{\mathcal{M}_2, g\}$ with $\Gamma^\lambda_{\mu\nu}, \nabla_\mu, \Upsilon_\mu, \mathcal{D}_\mu, R_{\alpha\beta\mu\nu}, C_{\mu\nu}$ defined in section 1.

Theorem 11. *On the pseudo-Riemannian space $\mathcal{V}_2$ there exists the solution $H \in \Lambda_1$, $B_\mu \in \Lambda_2 \mathbb{T}_1$ of the system of equations that consists of (21) and*

$$\mathcal{D}_\mu H = 0, \quad H^2 = 1. \quad (38)$$

Proof. We may take

$$\begin{aligned} b_{121} &= -(\partial_1 g_{22} g_{12} - \partial_2 g_{11} g_{22}) / (4 g_{22}) \\ b_{122} &= -(\partial_2 g_{22} g_{12} + \partial_1 g_{22} g_{22} - 2 \partial_2 g_{12} g_{22}) / (4 g_{22}) \end{aligned}$$

and check that $H = dx^1 / \sqrt{g^{11}}$ and $B_\mu = \frac{1}{2} b_{\alpha\beta\mu} dx^\alpha \wedge dx^\beta$ satisfy (21) and (38).

9 Addendum.

Here are formulas for the components $b_{\alpha\beta\mu}$ of B_μ . These B_μ together with the H, I, K from (14) satisfy equations (10-12)

$$\begin{aligned} b_{121} = & (\partial_1 g_{44} g_{14} g_{23} g_{24}^2 g_{33} g_{34} - \partial_1 g_{44} g_{13} g_{24}^3 g_{33} g_{34} - 2 \partial_1 g_{44} g_{14} g_{23}^2 g_{24} g_{34}^2 + \\ & 2 \partial_1 g_{44} g_{13} g_{23} g_{24}^2 g_{34}^2 + \partial_1 g_{44} g_{14} g_{22} g_{23} g_{34}^3 - \partial_1 g_{44} g_{13} g_{22} g_{24} g_{34}^3 + \partial_1 g_{44} g_{14} g_{23}^2 g_{24} g_{33} g_{44} - \\ & \partial_1 g_{44} g_{13} g_{23} g_{24}^2 g_{33} g_{44} - 2 \partial_1 g_{34} g_{14} g_{23} g_{24}^2 g_{33} g_{44} + 2 \partial_1 g_{34} g_{13} g_{24}^3 g_{33} g_{44} - \\ & \partial_1 g_{44} g_{14} g_{22} g_{24} g_{33}^2 g_{44} + \partial_1 g_{44} g_{12} g_{24}^2 g_{33}^2 g_{44} + 2 \partial_1 g_{34} g_{14} g_{23}^2 g_{24} g_{34} g_{44} - 2 \partial_1 g_{34} g_{13} g_{23} g_{24}^2 g_{34} g_{44} + \\ & \partial_1 g_{33} g_{14} g_{23} g_{24}^2 g_{34} g_{44} - \partial_1 g_{33} g_{13} g_{24}^3 g_{34} g_{44} + 2 \partial_1 g_{44} g_{13} g_{22} g_{24} g_{33} g_{34} g_{44} + \\ & 2 \partial_1 g_{34} g_{14} g_{22} g_{24} g_{33} g_{34} g_{44} - 2 \partial_1 g_{44} g_{12} g_{23} g_{24} g_{33} g_{34} g_{44} - 2 \partial_1 g_{34} g_{12} g_{24}^2 g_{33} g_{34} g_{44} - \\ & \partial_1 g_{44} g_{13} g_{22} g_{23} g_{34}^2 g_{44} - 2 \partial_1 g_{34} g_{14} g_{22} g_{23} g_{34}^2 g_{44} + \partial_1 g_{44} g_{12} g_{23}^2 g_{34}^2 g_{44} + 2 \partial_1 g_{24} g_{14} g_{23}^2 g_{34}^2 g_{44} - \\ & \partial_1 g_{33} g_{14} g_{22} g_{24} g_{34}^2 g_{44} + 2 \partial_1 g_{34} g_{12} g_{23} g_{24} g_{34}^2 g_{44} - 2 \partial_1 g_{24} g_{13} g_{23} g_{24} g_{34}^2 g_{44} - \\ & 2 \partial_1 g_{23} g_{14} g_{23} g_{24} g_{34}^2 g_{44} + \partial_1 g_{33} g_{12} g_{24}^2 g_{34}^2 g_{44} + 2 \partial_1 g_{23} g_{13} g_{24}^2 g_{34}^2 g_{44} - 2 \partial_1 g_{24} g_{14} g_{22} g_{33} g_{34}^2 g_{44} + \\ & 2 \partial_1 g_{24} g_{12} g_{24} g_{33} g_{34}^2 g_{44} + \partial_1 g_{22} g_{14} g_{24} g_{33} g_{34}^2 g_{44} - \partial_2 g_{11} g_{24}^2 g_{33} g_{34}^2 g_{44} + 2 \partial_1 g_{24} g_{13} g_{22} g_{34}^3 g_{44} + \\ & 2 \partial_1 g_{23} g_{14} g_{22} g_{34}^3 g_{44} - 2 \partial_1 g_{24} g_{12} g_{23} g_{34}^3 g_{44} - \partial_1 g_{22} g_{14} g_{23} g_{34}^3 g_{44} - 2 \partial_1 g_{23} g_{12} g_{24} g_{34}^3 g_{44} - \\ & \partial_1 g_{22} g_{13} g_{24} g_{34}^3 g_{44} + 2 \partial_2 g_{11} g_{23} g_{24} g_{34}^3 g_{44} + \partial_1 g_{22} g_{12} g_{34}^4 g_{44} - \partial_2 g_{11} g_{22} g_{34}^4 g_{44} - \\ & \partial_1 g_{33} g_{14} g_{23}^2 g_{24} g_{44}^2 + \partial_1 g_{33} g_{13} g_{23} g_{24}^2 g_{44}^2 - 2 \partial_1 g_{24} g_{14} g_{23}^2 g_{33} g_{44}^2 - 2 \partial_1 g_{34} g_{13} g_{22} g_{24} g_{33} g_{44}^2 + \\ & 2 \partial_1 g_{34} g_{12} g_{23} g_{24} g_{33} g_{44}^2 + 2 \partial_1 g_{24} g_{13} g_{23} g_{24} g_{33} g_{44}^2 + 2 \partial_1 g_{23} g_{14} g_{23} g_{24} g_{33} g_{44}^2 - \end{aligned}$$

$$\begin{aligned}
& 2 \partial_1 g_{23} g_{13} g_{24}^2 g_{33} g_{44}^2 + 2 \partial_1 g_{24} g_{14} g_{22} g_{33}^2 g_{44}^2 - 2 \partial_1 g_{24} g_{12} g_{24} g_{33}^2 g_{44}^2 - \partial_1 g_{22} g_{14} g_{24} g_{33}^2 g_{44}^2 + \\
& \partial_2 g_{11} g_{24}^2 g_{33}^2 g_{44}^2 + 2 \partial_1 g_{34} g_{13} g_{22} g_{23} g_{34} g_{44}^2 + \partial_1 g_{33} g_{14} g_{22} g_{23} g_{34} g_{44}^2 - 2 \partial_1 g_{34} g_{12} g_{23}^2 g_{34} g_{44}^2 + \\
& \partial_1 g_{33} g_{13} g_{22} g_{24} g_{34} g_{44}^2 - 2 \partial_1 g_{33} g_{12} g_{23} g_{24} g_{34} g_{44}^2 - 2 \partial_1 g_{24} g_{13} g_{22} g_{33} g_{34} g_{44}^2 - \\
& 2 \partial_1 g_{23} g_{14} g_{22} g_{33} g_{34} g_{44}^2 + 2 \partial_1 g_{24} g_{12} g_{23} g_{33} g_{34} g_{44}^2 + \partial_1 g_{22} g_{14} g_{23} g_{33} g_{34} g_{44}^2 + \\
& 2 \partial_1 g_{23} g_{12} g_{24} g_{33} g_{34} g_{44}^2 + \partial_1 g_{22} g_{13} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_2 g_{11} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - \\
& 2 \partial_1 g_{23} g_{13} g_{22} g_{34}^2 g_{44}^2 + 2 \partial_1 g_{23} g_{12} g_{23} g_{34}^2 g_{44}^2 + \partial_1 g_{22} g_{13} g_{23} g_{34}^2 g_{44}^2 - \partial_2 g_{11} g_{23}^2 g_{34}^2 g_{44}^2 - \\
& 2 \partial_1 g_{22} g_{12} g_{33} g_{34}^2 g_{44}^2 + 2 \partial_2 g_{11} g_{22} g_{33} g_{34}^2 g_{44}^2 - \partial_1 g_{33} g_{13} g_{22} g_{23} g_{44}^3 + \partial_1 g_{33} g_{12} g_{23}^2 g_{44}^3 + \\
& 2 \partial_1 g_{23} g_{13} g_{22} g_{33} g_{44}^3 - 2 \partial_1 g_{23} g_{12} g_{23} g_{33} g_{44}^3 - \partial_1 g_{22} g_{13} g_{23} g_{33} g_{44}^3 + \partial_2 g_{11} g_{23}^2 g_{33} g_{44}^3 + \\
& \partial_1 g_{22} g_{12} g_{33}^2 g_{44}^3 - \partial_2 g_{11} g_{22} g_{33}^2 g_{44}^3) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44}) (g_{24}^2 g_{33} - 2 g_{23} g_{24} g_{34} + \\
& g_{22} g_{34}^2 + g_{23}^2 g_{44} - g_{22} g_{33} g_{44})) \\
& b_{122} = (\partial_2 g_{44} g_{14} g_{23} g_{24}^2 g_{33} g_{34} - \partial_2 g_{44} g_{13} g_{24}^3 g_{33} g_{34} - 2 \partial_2 g_{44} g_{14} g_{23}^2 g_{24} g_{34}^2 + \\
& 2 \partial_2 g_{44} g_{13} g_{23} g_{24}^2 g_{34}^2 + \partial_2 g_{44} g_{14} g_{22} g_{23} g_{34}^3 - \partial_2 g_{44} g_{13} g_{22} g_{24} g_{34}^3 + \partial_2 g_{44} g_{14} g_{23}^2 g_{24} g_{33} g_{44} - \\
& \partial_2 g_{44} g_{13} g_{23} g_{24}^2 g_{33} g_{44} - 2 \partial_2 g_{34} g_{14} g_{23} g_{24}^2 g_{33} g_{44} + 2 \partial_2 g_{34} g_{13} g_{24}^3 g_{33} g_{44} - \\
& \partial_2 g_{44} g_{14} g_{22} g_{24} g_{33}^2 g_{44} + \partial_2 g_{44} g_{12} g_{24}^2 g_{33}^2 g_{44} + 2 \partial_2 g_{34} g_{14} g_{23}^2 g_{24} g_{34} g_{44} - 2 \partial_2 g_{34} g_{13} g_{23} g_{24}^2 g_{34} g_{44} + \\
& \partial_2 g_{33} g_{14} g_{23} g_{24}^2 g_{34} g_{44} - \partial_2 g_{33} g_{13} g_{24}^2 g_{34} g_{44} + 2 \partial_2 g_{44} g_{13} g_{22} g_{24} g_{33} g_{34} g_{44} + \\
& 2 \partial_2 g_{34} g_{14} g_{22} g_{24} g_{33} g_{34} g_{44} - 2 \partial_2 g_{44} g_{12} g_{23} g_{24} g_{33} g_{34} g_{44} - 2 \partial_2 g_{34} g_{12} g_{24}^2 g_{33} g_{34} g_{44} - \\
& \partial_2 g_{44} g_{13} g_{22} g_{23} g_{34}^2 g_{44} - 2 \partial_2 g_{34} g_{14} g_{22} g_{23} g_{34}^2 g_{44} + \partial_2 g_{44} g_{12} g_{23}^2 g_{34}^2 g_{44} + 2 \partial_2 g_{24} g_{14} g_{23}^2 g_{34}^2 g_{44} - \\
& \partial_2 g_{33} g_{14} g_{22} g_{24} g_{34}^2 g_{44} + 2 \partial_2 g_{34} g_{12} g_{23} g_{24} g_{34}^2 g_{44} - 2 \partial_2 g_{24} g_{13} g_{23} g_{24} g_{34}^2 g_{44} - \\
& 2 \partial_2 g_{23} g_{14} g_{23} g_{24} g_{34}^2 g_{44} + \partial_2 g_{33} g_{12} g_{24}^2 g_{34}^2 g_{44} + 2 \partial_2 g_{23} g_{13} g_{24}^2 g_{34}^2 g_{44} - 2 \partial_2 g_{24} g_{14} g_{22} g_{33} g_{34}^2 g_{44} + \\
& 2 \partial_2 g_{24} g_{12} g_{24} g_{33} g_{34}^2 g_{44} + \partial_2 g_{22} g_{14} g_{24} g_{33} g_{34}^2 g_{44} + \partial_1 g_{22} g_{24}^2 g_{33} g_{34}^2 g_{44} - 2 \partial_2 g_{12} g_{24}^2 g_{33} g_{34}^2 g_{44} + \\
& 2 \partial_2 g_{24} g_{13} g_{22} g_{34}^3 g_{44} + 2 \partial_2 g_{23} g_{14} g_{22} g_{34}^3 g_{44} - 2 \partial_2 g_{24} g_{12} g_{23} g_{34}^3 g_{44} - \partial_2 g_{22} g_{14} g_{23} g_{34}^3 g_{44} - \\
& 2 \partial_2 g_{23} g_{12} g_{24} g_{34}^3 g_{44} - \partial_2 g_{22} g_{13} g_{24} g_{34}^3 g_{44} - 2 \partial_1 g_{22} g_{23} g_{24} g_{34}^3 g_{44} + 4 \partial_2 g_{12} g_{23} g_{24} g_{34}^3 g_{44} + \\
& \partial_2 g_{22} g_{12} g_{34}^4 g_{44} + \partial_1 g_{22} g_{22} g_{34}^4 g_{44} - 2 \partial_2 g_{12} g_{22} g_{34}^4 g_{44} - \partial_2 g_{33} g_{14} g_{23}^2 g_{24} g_{44}^2 + \\
& \partial_2 g_{33} g_{13} g_{23} g_{24}^2 g_{44}^2 - 2 \partial_2 g_{24} g_{14} g_{23}^2 g_{33} g_{44}^2 - 2 \partial_2 g_{34} g_{13} g_{22} g_{24} g_{33} g_{44}^2 + 2 \partial_2 g_{34} g_{12} g_{23} g_{24} g_{33} g_{44}^2 + \\
& 2 \partial_2 g_{24} g_{13} g_{23} g_{24} g_{33} g_{44}^2 + 2 \partial_2 g_{23} g_{14} g_{23} g_{24} g_{33} g_{44}^2 - 2 \partial_2 g_{23} g_{13} g_{24}^2 g_{33} g_{44}^2 + \\
& 2 \partial_2 g_{24} g_{14} g_{22} g_{33}^2 g_{44}^2 - 2 \partial_2 g_{24} g_{12} g_{24} g_{33}^2 g_{44}^2 - \partial_2 g_{22} g_{14} g_{24} g_{33}^2 g_{44}^2 - \partial_1 g_{22} g_{24}^2 g_{33}^2 g_{44}^2 + \\
& 2 \partial_2 g_{12} g_{24}^2 g_{33}^2 g_{44}^2 + 2 \partial_2 g_{34} g_{13} g_{22} g_{23} g_{34} g_{44}^2 + \partial_2 g_{33} g_{14} g_{22} g_{23} g_{34} g_{44}^2 - 2 \partial_2 g_{34} g_{12} g_{23}^2 g_{34} g_{44}^2 + \\
& \partial_2 g_{33} g_{13} g_{22} g_{24} g_{34} g_{44}^2 - 2 \partial_2 g_{33} g_{12} g_{23} g_{24} g_{34} g_{44}^2 - 2 \partial_2 g_{24} g_{13} g_{22} g_{33} g_{34} g_{44}^2 - \\
& 2 \partial_2 g_{23} g_{14} g_{22} g_{33} g_{34} g_{44}^2 + 2 \partial_2 g_{24} g_{12} g_{23} g_{33} g_{34} g_{44}^2 + \partial_2 g_{22} g_{14} g_{23} g_{33} g_{34} g_{44}^2 + \\
& 2 \partial_2 g_{23} g_{12} g_{24} g_{33} g_{34} g_{44}^2 + \partial_2 g_{22} g_{13} g_{24} g_{33} g_{34} g_{44}^2 + 2 \partial_1 g_{22} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - \\
& 4 \partial_2 g_{12} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_2 g_{23} g_{13} g_{22} g_{34}^2 g_{44}^2 + 2 \partial_2 g_{23} g_{12} g_{23} g_{34}^2 g_{44}^2 + \partial_2 g_{22} g_{13} g_{23} g_{34}^2 g_{44}^2 + \\
& \partial_1 g_{22} g_{23}^2 g_{34}^2 g_{44}^2 - 2 \partial_2 g_{12} g_{23}^2 g_{34}^2 g_{44}^2 - 2 \partial_2 g_{22} g_{12} g_{33} g_{34}^2 g_{44}^2 - 2 \partial_1 g_{22} g_{22} g_{33} g_{34}^2 g_{44}^2 + \\
& 4 \partial_2 g_{12} g_{22} g_{33} g_{34}^2 g_{44}^2 - \partial_2 g_{33} g_{13} g_{22} g_{23} g_{44}^3 + \partial_2 g_{33} g_{12} g_{23}^2 g_{44}^3 + 2 \partial_2 g_{23} g_{13} g_{22} g_{33} g_{44}^3 - \\
& 2 \partial_2 g_{23} g_{12} g_{23} g_{33} g_{44}^3 - \partial_2 g_{22} g_{13} g_{23} g_{33} g_{44}^3 - \partial_1 g_{22} g_{23}^2 g_{33} g_{44}^3 + 2 \partial_2 g_{12} g_{23}^2 g_{33} g_{44}^3 + \\
& \partial_2 g_{22} g_{12} g_{33}^2 g_{44}^3 + \partial_1 g_{22} g_{22} g_{33}^2 g_{44}^3 - 2 \partial_2 g_{12} g_{22} g_{33}^2 g_{44}^3) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44}) (g_{24}^2 g_{33} - \\
& 2 g_{23} g_{24} g_{34} + g_{22} g_{34}^2 + g_{23}^2 g_{44} - g_{22} g_{33} g_{44})) \\
& b_{123} = (\partial_3 g_{44} g_{14} g_{23} g_{24}^2 g_{33} g_{34} - \partial_3 g_{44} g_{13} g_{24}^3 g_{33} g_{34} - 2 \partial_3 g_{44} g_{14} g_{23}^2 g_{24} g_{34}^2 +
\end{aligned}$$

$$\begin{aligned}
& 2 \partial_3 g_{44} g_{13} g_{23} g_{24}^2 g_{34}^2 + \partial_3 g_{44} g_{14} g_{22} g_{23} g_{34}^3 - \partial_3 g_{44} g_{13} g_{22} g_{24} g_{34}^3 + \partial_3 g_{44} g_{14} g_{23}^2 g_{24} g_{33} g_{44} - \\
& \partial_3 g_{44} g_{13} g_{23} g_{24}^2 g_{33} g_{44} - 2 \partial_3 g_{34} g_{14} g_{23} g_{24}^2 g_{33} g_{44} + 2 \partial_3 g_{34} g_{13} g_{24}^2 g_{33} g_{44} - \\
& \partial_3 g_{44} g_{14} g_{22} g_{24} g_{33}^2 g_{44} + \partial_3 g_{44} g_{12} g_{24}^2 g_{33}^2 g_{44} + 2 \partial_3 g_{34} g_{14} g_{23}^2 g_{24} g_{34} g_{44} - 2 \partial_3 g_{34} g_{13} g_{23} g_{24}^2 g_{34} g_{44} + \\
& \partial_3 g_{33} g_{14} g_{23} g_{24}^2 g_{34} g_{44} - \partial_3 g_{33} g_{13} g_{24}^3 g_{34} g_{44} + 2 \partial_3 g_{44} g_{13} g_{22} g_{24} g_{33} g_{34} g_{44} + \\
& 2 \partial_3 g_{34} g_{14} g_{22} g_{24} g_{33} g_{34} g_{44} - 2 \partial_3 g_{44} g_{12} g_{23} g_{24} g_{33} g_{34} g_{44} - 2 \partial_3 g_{34} g_{12} g_{24}^2 g_{33} g_{34} g_{44} - \\
& \partial_3 g_{44} g_{13} g_{22} g_{23} g_{34}^2 g_{44} - 2 \partial_3 g_{34} g_{14} g_{22} g_{23} g_{34}^2 g_{44} + \partial_3 g_{44} g_{12} g_{23}^2 g_{34}^2 g_{44} + 2 \partial_3 g_{24} g_{14} g_{23}^2 g_{34}^2 g_{44} - \\
& \partial_3 g_{33} g_{14} g_{22} g_{24} g_{34}^2 g_{44} + 2 \partial_3 g_{34} g_{12} g_{23} g_{24} g_{34}^2 g_{44} - 2 \partial_3 g_{24} g_{13} g_{23} g_{24} g_{34}^2 g_{44} - \\
& 2 \partial_3 g_{23} g_{14} g_{23} g_{24} g_{34}^2 g_{44} + \partial_3 g_{33} g_{12} g_{24}^2 g_{34}^2 g_{44} + 2 \partial_3 g_{23} g_{13} g_{24}^2 g_{34}^2 g_{44} - 2 \partial_3 g_{24} g_{14} g_{22} g_{33} g_{34}^2 g_{44} + \\
& 2 \partial_3 g_{24} g_{12} g_{24} g_{33} g_{34}^2 g_{44} + \partial_3 g_{22} g_{14} g_{24} g_{33} g_{34}^2 g_{44} + \partial_1 g_{23} g_{24}^2 g_{33} g_{34}^2 g_{44} - \partial_2 g_{13} g_{24}^2 g_{33} g_{34}^2 g_{44} - \\
& \partial_3 g_{12} g_{24}^2 g_{33} g_{34}^2 g_{44} + 2 \partial_3 g_{24} g_{13} g_{22} g_{34}^3 g_{44} + 2 \partial_3 g_{23} g_{14} g_{22} g_{34}^3 g_{44} - 2 \partial_3 g_{24} g_{12} g_{23} g_{34}^3 g_{44} - \\
& \partial_3 g_{22} g_{14} g_{23} g_{34}^3 g_{44} - 2 \partial_3 g_{23} g_{12} g_{24} g_{34}^3 g_{44} - \partial_3 g_{22} g_{13} g_{24} g_{34}^3 g_{44} - 2 \partial_1 g_{23} g_{23} g_{24} g_{34}^3 g_{44} + \\
& 2 \partial_2 g_{13} g_{23} g_{24} g_{34}^3 g_{44} + 2 \partial_3 g_{12} g_{23} g_{24} g_{34}^3 g_{44} + \partial_3 g_{22} g_{12} g_{34}^4 g_{44} + \partial_1 g_{23} g_{22} g_{34}^4 g_{44} - \\
& \partial_2 g_{13} g_{22} g_{34}^4 g_{44} - \partial_3 g_{12} g_{22} g_{34}^4 g_{44} - \partial_3 g_{33} g_{14} g_{23}^2 g_{24} g_{44}^2 + \partial_3 g_{33} g_{13} g_{23} g_{24}^2 g_{44}^2 - \\
& 2 \partial_3 g_{24} g_{14} g_{23}^2 g_{33} g_{44}^2 - 2 \partial_3 g_{34} g_{13} g_{22} g_{24} g_{33} g_{44}^2 + 2 \partial_3 g_{34} g_{12} g_{23} g_{24} g_{33} g_{44}^2 + \\
& 2 \partial_3 g_{24} g_{13} g_{23} g_{24} g_{33} g_{44}^2 + 2 \partial_3 g_{23} g_{14} g_{23} g_{24} g_{33} g_{44}^2 - 2 \partial_3 g_{23} g_{13} g_{24}^2 g_{33} g_{44}^2 + \\
& 2 \partial_3 g_{24} g_{14} g_{22} g_{33}^2 g_{44}^2 - 2 \partial_3 g_{24} g_{12} g_{24} g_{33}^2 g_{44}^2 - \partial_3 g_{22} g_{14} g_{24} g_{33}^2 g_{44}^2 - \partial_1 g_{23} g_{24}^2 g_{33}^2 g_{44}^2 + \\
& \partial_2 g_{13} g_{24}^2 g_{33}^2 g_{44}^2 + \partial_3 g_{12} g_{24}^2 g_{33}^2 g_{44}^2 + 2 \partial_3 g_{34} g_{13} g_{22} g_{23} g_{34} g_{44}^2 + \partial_3 g_{33} g_{14} g_{22} g_{23} g_{34} g_{44}^2 - \\
& 2 \partial_3 g_{34} g_{12} g_{23}^2 g_{34} g_{44}^2 + \partial_3 g_{33} g_{13} g_{22} g_{24} g_{34} g_{44}^2 - 2 \partial_3 g_{33} g_{12} g_{23} g_{24} g_{34} g_{44}^2 - \\
& 2 \partial_3 g_{24} g_{13} g_{22} g_{33} g_{34} g_{44}^2 + 2 \partial_3 g_{23} g_{14} g_{22} g_{33} g_{34} g_{44}^2 + \partial_3 g_{22} g_{13} g_{24} g_{33} g_{34} g_{44}^2 + \\
& 2 \partial_1 g_{23} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_2 g_{13} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_3 g_{12} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - \\
& 2 \partial_3 g_{23} g_{13} g_{22} g_{34}^2 g_{44}^2 + 2 \partial_3 g_{23} g_{12} g_{23} g_{34}^2 g_{44}^2 + \partial_3 g_{22} g_{13} g_{23} g_{34}^2 g_{44}^2 + \partial_1 g_{23} g_{23}^2 g_{34}^2 g_{44}^2 - \\
& \partial_2 g_{13} g_{23}^2 g_{34}^2 g_{44}^2 - \partial_3 g_{12} g_{23}^2 g_{34}^2 g_{44}^2 - 2 \partial_3 g_{22} g_{12} g_{33} g_{34}^2 g_{44}^2 - 2 \partial_1 g_{23} g_{22} g_{33} g_{34}^2 g_{44}^2 + \\
& 2 \partial_2 g_{13} g_{22} g_{33} g_{34}^2 g_{44}^2 + 2 \partial_3 g_{12} g_{22} g_{33} g_{34}^2 g_{44}^2 - \partial_3 g_{33} g_{13} g_{22} g_{23} g_{34}^3 + \partial_3 g_{33} g_{12} g_{23}^2 g_{34}^3 + \\
& 2 \partial_3 g_{23} g_{13} g_{22} g_{33} g_{34}^3 - 2 \partial_3 g_{23} g_{12} g_{23} g_{33} g_{34}^3 - \partial_3 g_{22} g_{13} g_{23} g_{33} g_{34}^3 - \partial_1 g_{23} g_{23}^2 g_{33} g_{34}^3 + \\
& \partial_2 g_{13} g_{23}^2 g_{33} g_{34}^3 + \partial_3 g_{12} g_{23}^2 g_{33} g_{34}^3 + \partial_3 g_{22} g_{12} g_{33}^2 g_{44}^3 + \partial_1 g_{23} g_{22} g_{33}^2 g_{44}^3 - \\
& \partial_2 g_{13} g_{22} g_{33}^2 g_{44}^3 - \partial_3 g_{12} g_{22} g_{33}^2 g_{44}^3) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44}) (g_{24}^2 g_{33} - 2 g_{23} g_{24} g_{34} + \\
& g_{22} g_{34}^2 + g_{23}^2 g_{44} - g_{22} g_{33} g_{44}))
\end{aligned}$$

$$\begin{aligned}
b_{124} = & (\partial_4 g_{44} g_{14} g_{23} g_{24}^2 g_{33} g_{34} - \partial_4 g_{44} g_{13} g_{24}^3 g_{33} g_{34} - 2 \partial_4 g_{44} g_{14} g_{23}^2 g_{24} g_{34}^2 + \\
& 2 \partial_4 g_{44} g_{13} g_{23} g_{24}^2 g_{34}^2 + \partial_4 g_{44} g_{14} g_{22} g_{23} g_{34}^3 - \partial_4 g_{44} g_{13} g_{22} g_{24} g_{34}^3 + \partial_4 g_{44} g_{14} g_{23}^2 g_{24} g_{33} g_{44} - \\
& \partial_4 g_{44} g_{13} g_{23} g_{24}^2 g_{33} g_{44} - 2 \partial_4 g_{34} g_{14} g_{23} g_{24}^2 g_{33} g_{44} + 2 \partial_4 g_{34} g_{13} g_{24}^2 g_{33} g_{44} - \\
& \partial_4 g_{44} g_{14} g_{22} g_{24} g_{33}^2 g_{44} + \partial_4 g_{44} g_{12} g_{24}^2 g_{33}^2 g_{44} + 2 \partial_4 g_{34} g_{14} g_{23}^2 g_{24} g_{34} g_{44} - 2 \partial_4 g_{34} g_{13} g_{23} g_{24}^2 g_{34} g_{44} + \\
& \partial_4 g_{33} g_{14} g_{23} g_{24}^2 g_{34} g_{44} - \partial_4 g_{33} g_{13} g_{24}^3 g_{34} g_{44} + 2 \partial_4 g_{44} g_{13} g_{22} g_{24} g_{33} g_{34} g_{44} + \\
& 2 \partial_4 g_{34} g_{14} g_{22} g_{24} g_{33} g_{34} g_{44} - 2 \partial_4 g_{44} g_{12} g_{23} g_{24} g_{33} g_{34} g_{44} - 2 \partial_4 g_{34} g_{12} g_{24}^2 g_{33} g_{34} g_{44} - \\
& \partial_4 g_{44} g_{13} g_{22} g_{23} g_{34}^2 g_{44} - 2 \partial_4 g_{34} g_{14} g_{22} g_{23} g_{34}^2 g_{44} + \partial_4 g_{44} g_{12} g_{23}^2 g_{34}^2 g_{44} + 2 \partial_4 g_{24} g_{14} g_{23}^2 g_{34}^2 g_{44} - \\
& \partial_4 g_{33} g_{14} g_{22} g_{24} g_{34}^2 g_{44} + 2 \partial_4 g_{34} g_{12} g_{23} g_{24} g_{34}^2 g_{44} - 2 \partial_4 g_{24} g_{13} g_{23} g_{24} g_{34}^2 g_{44} - \\
& 2 \partial_4 g_{23} g_{14} g_{23} g_{24} g_{34}^2 g_{44} + \partial_4 g_{33} g_{12} g_{24}^2 g_{34}^2 g_{44} + 2 \partial_4 g_{23} g_{13} g_{24}^2 g_{34}^2 g_{44} - 2 \partial_4 g_{24} g_{14} g_{22} g_{33} g_{34}^2 g_{44} +
\end{aligned}$$

$$\begin{aligned}
& 2 \partial_4 g_{24} g_{12} g_{24} g_{33} g_{34}^2 g_{44} + \partial_4 g_{22} g_{14} g_{24} g_{33} g_{34}^2 g_{44} + \partial_1 g_{24} g_{24}^2 g_{33} g_{34}^2 g_{44} - \partial_2 g_{14} g_{24}^2 g_{33} g_{34}^2 g_{44} - \\
& \partial_4 g_{12} g_{24}^2 g_{33} g_{34}^2 g_{44} + 2 \partial_4 g_{24} g_{13} g_{22} g_{34}^3 g_{44} + 2 \partial_4 g_{23} g_{14} g_{22} g_{34}^3 g_{44} - 2 \partial_4 g_{24} g_{12} g_{23} g_{34}^3 g_{44} - \\
& \partial_4 g_{22} g_{14} g_{23} g_{34}^3 g_{44} - 2 \partial_4 g_{23} g_{12} g_{24} g_{34}^3 g_{44} - \partial_4 g_{22} g_{13} g_{24} g_{34}^3 g_{44} - 2 \partial_1 g_{24} g_{23} g_{24} g_{34}^3 g_{44} + \\
& 2 \partial_2 g_{14} g_{23} g_{24} g_{34}^3 g_{44} + 2 \partial_4 g_{12} g_{23} g_{24} g_{34}^3 g_{44} + \partial_4 g_{22} g_{12} g_{34}^4 g_{44} + \partial_1 g_{24} g_{22} g_{34}^4 g_{44} - \\
& \partial_2 g_{14} g_{22} g_{34}^4 g_{44} - \partial_4 g_{12} g_{22} g_{34}^4 g_{44} - \partial_4 g_{33} g_{14} g_{23}^2 g_{24} g_{44}^2 + \partial_4 g_{33} g_{13} g_{23} g_{24}^2 g_{44}^2 - \\
& 2 \partial_4 g_{24} g_{14} g_{23}^2 g_{33} g_{44}^2 - 2 \partial_4 g_{34} g_{13} g_{22} g_{24} g_{33} g_{44}^2 + 2 \partial_4 g_{34} g_{12} g_{23} g_{24} g_{33} g_{44}^2 + \\
& 2 \partial_4 g_{24} g_{13} g_{23} g_{24} g_{33} g_{44}^2 + 2 \partial_4 g_{23} g_{14} g_{23} g_{24} g_{33} g_{44}^2 - 2 \partial_4 g_{23} g_{13} g_{24}^2 g_{33} g_{44}^2 + \\
& 2 \partial_4 g_{24} g_{14} g_{22} g_{33}^2 g_{44}^2 - 2 \partial_4 g_{24} g_{12} g_{24} g_{33}^2 g_{44}^2 - \partial_4 g_{22} g_{14} g_{24} g_{33}^2 g_{44}^2 - \partial_1 g_{24} g_{24}^2 g_{33}^2 g_{44}^2 + \\
& \partial_2 g_{14} g_{24}^2 g_{33}^2 g_{44}^2 + \partial_4 g_{12} g_{24}^2 g_{33}^2 g_{44}^2 + 2 \partial_4 g_{34} g_{13} g_{22} g_{23} g_{34} g_{44}^2 + \partial_4 g_{33} g_{14} g_{22} g_{23} g_{34} g_{44}^2 - \\
& 2 \partial_4 g_{34} g_{12} g_{23}^2 g_{34} g_{44}^2 + \partial_4 g_{33} g_{13} g_{22} g_{24} g_{34} g_{44}^2 - 2 \partial_4 g_{33} g_{12} g_{23} g_{24} g_{34} g_{44}^2 - \\
& 2 \partial_4 g_{24} g_{13} g_{22} g_{33} g_{34} g_{44}^2 - 2 \partial_4 g_{23} g_{14} g_{22} g_{33} g_{34} g_{44}^2 + 2 \partial_4 g_{24} g_{12} g_{23} g_{33} g_{34} g_{44}^2 + \\
& \partial_4 g_{22} g_{14} g_{23} g_{33} g_{34} g_{44}^2 + 2 \partial_4 g_{23} g_{12} g_{24} g_{33} g_{34} g_{44}^2 + \partial_4 g_{22} g_{13} g_{24} g_{33} g_{34} g_{44}^2 + \\
& 2 \partial_1 g_{24} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_2 g_{14} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - 2 \partial_4 g_{12} g_{23} g_{24} g_{33} g_{34} g_{44}^2 - \\
& 2 \partial_4 g_{23} g_{13} g_{22} g_{34}^2 g_{44}^2 + 2 \partial_4 g_{23} g_{12} g_{23} g_{34}^2 g_{44}^2 + \partial_4 g_{22} g_{13} g_{23} g_{34}^2 g_{44}^2 + \partial_1 g_{24} g_{23}^2 g_{34}^2 g_{44}^2 - \\
& \partial_2 g_{14} g_{23}^2 g_{34}^2 g_{44}^2 - \partial_4 g_{12} g_{23}^2 g_{34}^2 g_{44}^2 - 2 \partial_4 g_{22} g_{12} g_{33} g_{34}^2 g_{44}^2 - 2 \partial_1 g_{24} g_{22} g_{33} g_{34}^2 g_{44}^2 + \\
& 2 \partial_2 g_{14} g_{22} g_{33} g_{34}^2 g_{44}^2 + 2 \partial_4 g_{12} g_{22} g_{33} g_{34}^2 g_{44}^2 - \partial_4 g_{33} g_{13} g_{22} g_{23} g_{44}^3 + \partial_4 g_{33} g_{12} g_{23}^2 g_{44}^3 + \\
& 2 \partial_4 g_{23} g_{13} g_{22} g_{33} g_{44}^3 - 2 \partial_4 g_{23} g_{12} g_{23} g_{33} g_{44}^3 - \partial_4 g_{22} g_{13} g_{23} g_{33} g_{44}^3 - \partial_1 g_{24} g_{23}^2 g_{33} g_{44}^3 + \\
& \partial_2 g_{14} g_{23}^2 g_{33} g_{44}^3 + \partial_4 g_{12} g_{23}^2 g_{33} g_{44}^3 + \partial_4 g_{22} g_{12} g_{33}^2 g_{44}^3 + \partial_1 g_{24} g_{22} g_{33}^2 g_{44}^3 - \\
& \partial_2 g_{14} g_{22} g_{33}^2 g_{44}^3 - \partial_4 g_{12} g_{22} g_{33}^2 g_{44}^3) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44}) (g_{24}^2 g_{33} - 2 g_{23} g_{24} g_{34} + \\
& g_{22} g_{34}^2 + g_{23}^2 g_{44} - g_{22} g_{33} g_{44})) \\
b_{131} &= (\partial_1 g_{44} g_{14} g_{33} g_{34} - \partial_1 g_{44} g_{13} g_{34}^2 - 2 \partial_1 g_{34} g_{14} g_{33} g_{44} + 2 \partial_1 g_{34} g_{13} g_{34} g_{44} + \\
& \partial_1 g_{33} g_{14} g_{34} g_{44} - \partial_3 g_{11} g_{34}^2 g_{44} - \partial_1 g_{33} g_{13} g_{44}^2 + \partial_3 g_{11} g_{33} g_{44}^2) / (4 g_{44} (-g_{34}^2 + \\
& g_{33} g_{44})) \\
b_{132} &= (\partial_2 g_{44} g_{14} g_{33} g_{34} - \partial_2 g_{44} g_{13} g_{34}^2 - 2 \partial_2 g_{34} g_{14} g_{33} g_{44} + 2 \partial_2 g_{34} g_{13} g_{34} g_{44} + \\
& \partial_2 g_{33} g_{14} g_{34} g_{44} + \partial_1 g_{23} g_{34}^2 g_{44} - \partial_2 g_{13} g_{34}^2 g_{44} - \partial_3 g_{12} g_{34}^2 g_{44} - \partial_2 g_{33} g_{13} g_{44}^2 - \\
& \partial_1 g_{23} g_{33} g_{44}^2 + \partial_2 g_{13} g_{33} g_{44}^2 + \partial_3 g_{12} g_{33} g_{44}^2) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44})) \\
b_{133} &= (\partial_3 g_{44} g_{14} g_{33} g_{34} - \partial_3 g_{44} g_{13} g_{34}^2 - 2 \partial_3 g_{34} g_{14} g_{33} g_{44} + 2 \partial_3 g_{34} g_{13} g_{34} g_{44} + \\
& \partial_3 g_{33} g_{14} g_{34} g_{44} + \partial_1 g_{33} g_{34}^2 g_{44} - 2 \partial_3 g_{13} g_{34}^2 g_{44} - \partial_3 g_{33} g_{13} g_{44}^2 - \partial_1 g_{33} g_{33} g_{44}^2 + \\
& 2 \partial_3 g_{13} g_{33} g_{44}^2) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44})) \\
b_{134} &= (\partial_4 g_{44} g_{14} g_{33} g_{34} - \partial_4 g_{44} g_{13} g_{34}^2 - 2 \partial_4 g_{34} g_{14} g_{33} g_{44} + 2 \partial_4 g_{34} g_{13} g_{34} g_{44} + \\
& \partial_4 g_{33} g_{14} g_{34} g_{44} + \partial_1 g_{34} g_{34}^2 g_{44} - \partial_3 g_{14} g_{34}^2 g_{44} - \partial_4 g_{13} g_{34}^2 g_{44} - \partial_4 g_{33} g_{13} g_{44}^2 - \\
& \partial_1 g_{34} g_{33} g_{44}^2 + \partial_3 g_{14} g_{33} g_{44}^2 + \partial_4 g_{13} g_{33} g_{44}^2) / (4 g_{44} (-g_{34}^2 + g_{33} g_{44})) \\
b_{141} &= (-\partial_1 g_{44} g_{14}) / (4 g_{44}) \\
b_{142} &= (-\partial_2 g_{44} g_{14}) - \partial_1 g_{24} g_{44} + \partial_2 g_{14} g_{44} + \partial_4 g_{12} g_{44}) / (4 g_{44}) \\
b_{143} &= (-\partial_3 g_{44} g_{14}) - \partial_1 g_{34} g_{44} + \partial_3 g_{14} g_{44} + \partial_4 g_{13} g_{44}) / (4 g_{44}) \\
b_{144} &= (-\partial_4 g_{44} g_{14}) - \partial_1 g_{44} g_{44} + 2 \partial_4 g_{14} g_{44}) / (4 g_{44}) \\
b_{231} &= (-\partial_1 g_{44} g_{24} g_{33} g_{34}) + \partial_1 g_{44} g_{23} g_{34}^2 + 2 \partial_1 g_{34} g_{24} g_{33} g_{44} - 2 \partial_1 g_{34} g_{23} g_{34} g_{44} -
\end{aligned}$$

$$\begin{aligned}
& \partial_1 g_{33} g_{24} g_{34} g_{44} + \partial_1 g_{23} g_{34}^2 g_{44} - \partial_2 g_{13} g_{34}^2 g_{44} + \partial_3 g_{12} g_{34}^2 g_{44} + \partial_1 g_{33} g_{23} g_{44}^2 - \\
& \partial_1 g_{23} g_{33} g_{44}^2 + \partial_2 g_{13} g_{33} g_{44}^2 - \partial_3 g_{12} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44})) \\
b_{232} = & (- (\partial_2 g_{44} g_{24} g_{33} g_{34}) + \partial_2 g_{44} g_{23} g_{34}^2 + 2 \partial_2 g_{34} g_{24} g_{33} g_{44} - 2 \partial_2 g_{34} g_{23} g_{34} g_{44} - \\
& \partial_2 g_{33} g_{24} g_{34} g_{44} + \partial_3 g_{22} g_{34}^2 g_{44} + \partial_2 g_{33} g_{23} g_{44}^2 - \partial_3 g_{22} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44})) \\
b_{233} = & (- (\partial_3 g_{44} g_{24} g_{33} g_{34}) + \partial_3 g_{44} g_{23} g_{34}^2 + 2 \partial_3 g_{34} g_{24} g_{33} g_{44} - 2 \partial_3 g_{34} g_{23} g_{34} g_{44} - \\
& \partial_3 g_{33} g_{24} g_{34} g_{44} - \partial_2 g_{33} g_{34}^2 g_{44} + 2 \partial_3 g_{23} g_{34}^2 g_{44} + \partial_3 g_{33} g_{23} g_{44}^2 + \partial_2 g_{33} g_{33} g_{44}^2 - \\
& 2 \partial_3 g_{23} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44})) \\
b_{234} = & (- (\partial_4 g_{44} g_{24} g_{33} g_{34}) + \partial_4 g_{44} g_{23} g_{34}^2 + 2 \partial_4 g_{34} g_{24} g_{33} g_{44} - 2 \partial_4 g_{34} g_{23} g_{34} g_{44} - \\
& \partial_4 g_{33} g_{24} g_{34} g_{44} - \partial_2 g_{34} g_{34}^2 g_{44} + \partial_3 g_{24} g_{34}^2 g_{44} + \partial_4 g_{23} g_{34}^2 g_{44} + \partial_4 g_{33} g_{23} g_{44}^2 + \\
& \partial_2 g_{34} g_{33} g_{44}^2 - \partial_3 g_{24} g_{33} g_{44}^2 - \partial_4 g_{23} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44})) \\
b_{241} = & (- (\partial_1 g_{44} g_{24}) + \partial_1 g_{24} g_{44} - \partial_2 g_{14} g_{44} + \partial_4 g_{12} g_{44}) / (4 g_{44}) \\
b_{242} = & (- (\partial_2 g_{44} g_{24}) + \partial_4 g_{22} g_{44}) / (4 g_{44}) \\
b_{243} = & (- (\partial_3 g_{44} g_{24}) - \partial_2 g_{34} g_{44} + \partial_3 g_{24} g_{44} + \partial_4 g_{23} g_{44}) / (4 g_{44}) \\
b_{244} = & (- (\partial_4 g_{44} g_{24}) - \partial_2 g_{44} g_{44} + 2 \partial_4 g_{24} g_{44}) / (4 g_{44}) \\
b_{341} = & (- (\partial_1 g_{44} g_{34}) + \partial_1 g_{34} g_{44} - \partial_3 g_{14} g_{44} + \partial_4 g_{13} g_{44}) / (4 g_{44}) \\
b_{342} = & (- (\partial_2 g_{44} g_{34}) + \partial_2 g_{34} g_{44} - \partial_3 g_{24} g_{44} + \partial_4 g_{23} g_{44}) / (4 g_{44}) \\
b_{343} = & (- (\partial_3 g_{44} g_{34}) + \partial_4 g_{33} g_{44}) / (4 g_{44}) \\
b_{344} = & (- (\partial_4 g_{44} g_{34}) - \partial_3 g_{44} g_{44} + 2 \partial_4 g_{34} g_{44}) / (4 g_{44}).
\end{aligned}$$

Here are the formulas for $b_{\alpha\beta\mu}$ in the temporal gauge ($g_{11} = 1$, $g_{12} = g_{13} = g_{14} = 0$)

$$\begin{aligned}
b_{121} &= 0 \\
b_{122} &= -\partial_1 g_{22} / 4 \\
b_{123} &= -\partial_1 g_{23} / 4 \\
b_{124} &= -\partial_1 g_{24} / 4 \\
b_{131} &= 0 \\
b_{132} &= -\partial_1 g_{23} / 4 \\
b_{133} &= -\partial_1 g_{33} / 4 \\
b_{134} &= -\partial_1 g_{34} / 4 \\
b_{141} &= 0 \\
b_{142} &= -\partial_1 g_{24} / 4 \\
b_{143} &= -\partial_1 g_{34} / 4 \\
b_{144} &= -\partial_1 g_{44} / 4 \\
b_{231} &= -(\partial_1 g_{44} g_{24} g_{33} g_{34} - \partial_1 g_{44} g_{23} g_{34}^2 - 2 \partial_1 g_{34} g_{24} g_{33} g_{44} + 2 \partial_1 g_{34} g_{23} g_{34} g_{44} + \\
& \partial_1 g_{33} g_{24} g_{34} g_{44} - \partial_1 g_{23} g_{34}^2 g_{44} - \partial_1 g_{33} g_{23} g_{44}^2 + \partial_1 g_{23} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44})) \\
b_{232} &= -(\partial_2 g_{44} g_{24} g_{33} g_{34} - \partial_2 g_{44} g_{23} g_{34}^2 - 2 \partial_2 g_{34} g_{24} g_{33} g_{44} + 2 \partial_2 g_{34} g_{23} g_{34} g_{44} + \\
& \partial_2 g_{33} g_{24} g_{34} g_{44} - \partial_3 g_{22} g_{34}^2 g_{44} - \partial_2 g_{33} g_{23} g_{44}^2 + \partial_3 g_{22} g_{33} g_{44}^2) / (4 g_{44} (g_{34}^2 - g_{33} g_{44}))
\end{aligned}$$

$$\begin{aligned}
b_{233} &= -((\partial_3 g_{44} g_{24} g_{33} g_{34} - \partial_3 g_{44} g_{23} g_{34}^2 - 2 \partial_3 g_{34} g_{24} g_{33} g_{44} + 2 \partial_3 g_{34} g_{23} g_{34} g_{44} + \\
&\partial_3 g_{33} g_{24} g_{34} g_{44} + \partial_2 g_{33} g_{34}^2 g_{44} - 2 \partial_3 g_{23} g_{34}^2 g_{44} - \partial_3 g_{33} g_{23} g_{44}^2 - \partial_2 g_{33} g_{33} g_{44}^2 + \\
&2 \partial_3 g_{23} g_{33} g_{44}^2)/(4 g_{34}^2 g_{44} - 4 g_{33} g_{44}^2)) \\
b_{234} &= -((\partial_4 g_{44} g_{24} g_{33} g_{34} - \partial_4 g_{44} g_{23} g_{34}^2 - 2 \partial_4 g_{34} g_{24} g_{33} g_{44} + 2 \partial_4 g_{34} g_{23} g_{34} g_{44} + \\
&\partial_4 g_{33} g_{24} g_{34} g_{44} + \partial_2 g_{34} g_{34}^2 g_{44} - \partial_3 g_{24} g_{34}^2 g_{44} - \partial_4 g_{23} g_{34}^2 g_{44} - \partial_4 g_{33} g_{23} g_{44}^2 - \\
&\partial_2 g_{34} g_{33} g_{44}^2 + \partial_3 g_{24} g_{33} g_{44}^2 + \partial_4 g_{23} g_{33} g_{44}^2)/(4 g_{34}^2 g_{44} - 4 g_{33} g_{44}^2)) \\
b_{241} &= -(\partial_1 g_{44} g_{24}) + \partial_1 g_{24} g_{44})/(4 g_{44}) \\
b_{242} &= -(\partial_2 g_{44} g_{24}) + \partial_4 g_{22} g_{44})/(4 g_{44}) \\
b_{243} &= -(\partial_3 g_{44} g_{24}) - \partial_2 g_{34} g_{44} + \partial_3 g_{24} g_{44} + \partial_4 g_{23} g_{44})/(4 g_{44}) \\
b_{244} &= -(\partial_4 g_{44} g_{24}) - \partial_2 g_{44} g_{44} + 2 \partial_4 g_{24} g_{44})/(4 g_{44}) \\
b_{341} &= -(\partial_1 g_{44} g_{34} - \partial_1 g_{34} g_{44})/(4 g_{44}) \\
b_{342} &= -(\partial_2 g_{44} g_{34} - \partial_2 g_{34} g_{44} + \partial_3 g_{24} g_{44} - \partial_4 g_{23} g_{44})/(4 g_{44}) \\
b_{343} &= -(\partial_3 g_{44} g_{34} - \partial_4 g_{33} g_{44})/(4 g_{44}) \\
b_{344} &= -(\partial_4 g_{44} g_{34} + \partial_3 g_{44} g_{44} - 2 \partial_4 g_{34} g_{44})/(4 g_{44}).
\end{aligned}$$

Here are the formulas for $b_{\alpha\beta\mu}$ in case of the diagonal metric tensor $g_{\mu\nu} = \text{diag}(g_{11}, g_{22}, g_{33}, g_{44})$. These formulas are also valid for the space dimensions $n = 2, 3$ (see section 8)

$$b_{\alpha\beta\mu} = -\frac{1}{2}\partial_{[\alpha}g_{\beta]\mu}.$$

References

- [1] Marchuk N.G., Advances in Applied Clifford Algebras, v.8, N.1, (1998), p.181-225.(<http://xxx.lanl.gov/abs/math-ph/9811022>)
- [2] Marchuk N.G., Nuovo Cimento, 115B, 11, 2000.
- [3] Marchuk N.G., Dirac type tensor equations, to appear in Nuovo Cimento B in 2001.
- [4] Dirac P.A.M., Proc. Roy. Soc. Lond. A117 (1928) 610.
- [5] Ivanenko D., Landau L., Z. Phys., 48 (1928)340.
- [6] Kähler E., Rendiconti di Mat. (Roma) ser. 5, 21, (1962) 425.
- [7] Gürsey F., Nuovo Cimento, 3, p.988 (1956).
- [8] Hestenes D., *Space-Time Algebra*, Gordon and Breach, New York, 1966.

- [9] Hestenes D., J. Math. Phys., 8, pp.798-808 (1967).
- [10] Nakahara M., *Geometry, Topology and Physics*, Institute of Physics Publishing, Bristol and Philadelphia, 1998.